

BadgerCare Plus Benchmark Plan

Product List-Sorted by Generic Ingredient

The information provided below does not guarantee Wisconsin BadgerCare Plus Benchmark Plan coverage or payment. Real-time claim submission provides the most accurate member enrollment, drug coverage, and reimbursement determination. Please reference the date below when contacting Provider Services with questions about information received through this page. Providers may want to keep a copy of the page for their records.

| | | | | Benchmark Effective: October 1, 2011 | | | |
|-------------------------|-----------------|----------|-----|--------------------------------------|-------------------|----------|-----|
| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
| ACARBOSE | | | | ACETAMINOPHEN | | | |
| Oral | | | | Oral | | | |
| Tablet | | | | Suspension | | | |
| | 25 mg | 7/1/2009 | | | 650 mg/20.3 mL | 1/1/2011 | |
| | 50 mg | 7/1/2009 | | Tablet | | | |
| | 100 mg | 7/1/2009 | | | 325 mg | 2/1/2008 | |
| ACEBUTOLOL | | | | | 500 mg | 2/1/2008 | |
| Oral | | | | | 650 mg | 2/1/2008 | |
| Capsule | | | | Tablet ER | | | |
| | 200 mg | 2/1/2008 | | | 650 mg | 8/2/2010 | |
| | 400 mg | 2/1/2008 | | Rectal | | | |
| ACETAMINOPHEN | | | | Suppository | | | |
| Oral | | | | | 80 mg | 5/1/2008 | |
| Capsule | | | | | 120 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | 325 mg | 2/1/2008 | |
| Drops | | | | | 650 mg | 2/1/2008 | |
| | 100 mg/mL | 2/1/2008 | | ACETAMINOPHEN (Age 0-12) | | | |
| Drops Suspension | | | | Oral | | | |
| | 80 mg/0.8 mL | 2/1/2008 | | Tablet Chewable | | | |
| | 100 mg/mL | 2/1/2008 | | | 80 mg | 2/1/2008 | |
| Elixir | | | | Tablet Rapdis | | | |
| | 160 mg/5 mL | 2/1/2008 | | | 80 mg | 1/1/2011 | |
| Liquid | | | | | 160 mg | 1/1/2011 | |
| | 160 mg/5 mL | 2/1/2008 | | ACETAMINOPHEN and CODEINE | | | |
| | 500 mg/15 mL | 2/1/2008 | | Oral | | | |
| | 500 mg/5 mL | 1/1/2011 | | Elixir | | | |
| Solution | | | | | 120 mg-12 mg/5 mL | 2/1/2008 | |
| | 160 mg/5 mL | 2/1/2008 | | Tablet | | | |
| | 325 mg/10.15 mL | 2/1/2008 | | | 300 mg-15 mg | 2/1/2008 | |
| | 650 mg/20.3 mL | 2/1/2008 | | | 300 mg-30 mg | 2/1/2008 | |
| Suspension | | | | | 300 mg-60 mg | 2/1/2008 | |
| | 160 mg/5 mL | 2/1/2008 | | ACETAZOLAMIDE | | | |
| | 320 mg/10 mL | 1/1/2011 | | Oral | | | |
| | 325 mg/10.15 mL | 6/1/2008 | | Capsule ER | | | |
| | 640 mg/20 mL | 1/1/2011 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|----------------------|--------------------------------------------|----------|-----|--------------------------------------------|----------------------------------------|----------|-----|
| ACETAZOLAMIDE | | | | ACIDOPHILUS COMBINATIONS (Age 0-20) | | | |
| | Oral | | | | Oral | | |
| | Capsule ER | | | | Tablet | | |
| | 500 mg | 2/1/2009 | | | 25 million cell-100 mg | 1/1/2011 | |
| | | | | | 35 million cell-25 million cell | 1/1/2011 | |
| | ACETIC ACID | | | | Tablet Chewable | | |
| | Otic | | | | 7.5 billion cell-1 billion cell-500 mg | 1/1/2011 | |
| | Solution | | | | Tablet DR | | |
| | 2 % | 2/1/2008 | | | 250 mg-90 mg-350 mg-50 mg-150 mg | 1/1/2011 | |
| | ACETIC ACID and ALUMINUM | | | | | | |
| | Otic | | | | ACYCLOVIR | | |
| | Drops | | | | Oral | | |
| | 2 % | 2/1/2008 | | | Capsule | | |
| | | | | | 200 mg | 2/1/2008 | |
| | ACETYLCYSTEINE | | | | Suspension | | |
| | Miscell | | | | 200 mg/5 ml | 2/1/2008 | |
| | Vial | | | | Tablet | | |
| | 10 % (100 mg/mL) | 2/1/2008 | | | 400 mg | 2/1/2008 | |
| | 20 % (200 mg/mL) | 2/1/2008 | | | 800 mg | 2/1/2008 | |
| | ACIDOPHILUS and PECTIN (Age 0-20) | | | | ALBUTEROL | | |
| | Oral | | | | Inhalation | | |
| | Capsule | | | | Solution | | |
| | | 1/1/2011 | | | 5 mg/m; | 2/1/2008 | |
| | ACIDOPHILUS COMBINATIONS (Age 0-20) | | | | Vial-Neb | | |
| | Oral | | | | 1.25 mg/3 ml (Age 0-12) | 2/1/2008 | |
| | Capsule | | | | 2.5 mg/0.5 ml | 2/1/2008 | |
| | 10 billion cells (2 billion each) | 1/1/2011 | | | 2.5 mg/3 ml (0.083 %) | 2/1/2008 | |
| | 100 million cell | 1/1/2011 | | | Oral | | |
| | 16 billion cell | 1/1/2011 | | | Syrup | | |
| | 16 mg (1 billion cell) | 1/1/2011 | | | 2 mg/5 ml | 2/1/2008 | |
| | 175 mg | 1/1/2011 | | | Tablet | | |
| | 4 mg (1 billion cell) | 1/1/2011 | | | 2 mg | 2/1/2008 | |
| | 8 billion cell | 1/1/2011 | | | 4 mg | 2/1/2008 | |
| | Granuals | | | | Tablet (12hr) | | |
| | 100 million cell | 1/1/2011 | | | 4 mg | 2/1/2008 | |
| | Powder Pack | | | | 8 mg | 2/1/2008 | |
| | 7.5 billion cell-1 billion cell-500 mg | 1/1/2011 | | | ALENDRONATE | | |
| | Tablet | | | | Oral | | |
| | 1 billion cell-250 mg | 1/1/2011 | | | Tablet | | |
| | 1 million cell | 1/1/2011 | | | 5 mg | 2/1/2009 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----------------------------------------------|--------------------------|----------|-----|------------------------------------------|------------|----------|-----|
| ALENDRONATE | | | | AMANTADINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Syrup | | |
| | 10 mg | 2/1/2009 | | | 50 mg/5 mL | 2/1/2008 | |
| | 35 mg | 2/1/2009 | | | Tablet | | |
| | 40 mg | 2/1/2009 | | | 100 mg | 2/1/2008 | |
| | 70 mg | 2/1/2009 | | | | | |
| ALLOPURINOL | | | | AMILORIDE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 100 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| | 300 mg | 2/1/2008 | | | | | |
| ALPRAZOLAM | | | | AMILORIDE and HYDROCHLOROTHIAZIDE | | | |
| | Oral | | | | Oral | | |
| | Concentrate | | | | Tablet | | |
| | 1 mg/mL | 2/1/2008 | | | 5 mg-50 mg | 2/1/2008 | |
| | Tablet | | | | | | |
| | 0.25 mg | 2/1/2008 | | | | | |
| | 0.5 mg | 2/1/2008 | | | | | |
| | 1 mg | 2/1/2008 | | | | | |
| | 2 mg | 2/1/2008 | | | | | |
| | Tablet (24hr) | | | | | | |
| | 0.5 mg | 2/1/2008 | | | | | |
| | 1 mg | 2/1/2008 | | | | | |
| | 2 mg | 2/1/2008 | | | | | |
| | 3 mg | 2/1/2008 | | | | | |
| ALUMINUM and MAGNESIUM and SIMETHICONE | | | | AMINOCAPROIC ACID | | | |
| | Oral | | | | Oral | | |
| | Suspension | | | | Solution | | |
| | 200 mg-200 mg-20 mg/5 mL | 7/1/2011 | | | 250 mg/mL | 1/1/2010 | |
| | 400 mg-400 mg-40 mg/5 mL | 7/1/2011 | | | Tablet | | |
| | | | | | 500 mg | 1/1/2010 | |
| ALUMINUM HYDROXIDE | | | | AMINOPHYLLINE | | | |
| | Oral | | | | Oral | | |
| | Suspension | | | | Tablet | | |
| | 320 mg/5 mL | 2/1/2008 | | | 100 mg | 2/1/2008 | |
| | 600 mg/5 mL | 2/1/2008 | | | 200 mg | 2/1/2008 | |
| AMANTADINE | | | | AMIODARONE | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Tablet | | |
| | 100 mg | 2/1/2008 | | | 100 mg | 2/1/2008 | |
| | | | | | 200 mg | 2/1/2008 | |
| | | | | | 400 mg | 2/1/2008 | |
| N/D; N=New, D=Delete | | | | AMITRIPTYLINE | | | |
| | | | | | Oral | | |
| | | | | | Tablet | | |
| | | | | | 10 mg | 2/1/2008 | |
| | | | | | 25 mg | 2/1/2008 | |
| | | | | | 50 mg | 2/1/2008 | |
| | | | | | 75 mg | 2/1/2008 | |
| | | | | | 100 mg | 2/1/2008 | |
| | | | | | 150 mg | 2/1/2008 | |

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------------------------|---------------------|-----------|-----|-----------------------------------------------|-------------|-----------|-----|
| AMLODIPINE | | | | AMPHET ASP and AMPHET and D-AMPHET | | | |
| Oral | | | | Oral | | | |
| Tablet | | | | Capsule (24hr) | | | |
| | 2.5 mg | 4/2/2008 | | | 10 mg | 10/1/2010 | |
| | 5 mg | 4/2/2008 | | | 15 mg | 10/1/2010 | |
| | 10 mg | 4/2/2008 | | | 20 mg | 10/1/2010 | |
| | | | | | 25 mg | 10/1/2010 | |
| | | | | | 30 mg | 10/1/2010 | |
| AMOX TR-POTASSIUM CLAVULANATE | | | | Tablet | | | |
| Oral | | | | | 5 mg | 2/1/2008 | |
| Suspension Recon | | | | | 7.5 mg | 2/1/2008 | |
| | 200 mg-28.5 mg/5 mL | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | 250 mg-62.5 mg/5 mL | 10/1/2010 | | | 12.5 mg | 2/1/2008 | |
| | 400 mg-57 mg/5 mL | 2/1/2008 | | | 15 mg | 2/1/2008 | |
| | 600 mg-42.9 mg/5 mL | 2/1/2008 | | | 20 mg | 2/1/2008 | |
| | | | | | 30 mg | 2/1/2008 | |
| Tab Chew | | | | AMPICILLIN | | | |
| | 200 mg-28.5 mg | 2/1/2008 | | Oral | | | |
| | 400 mg-57 mg | 2/1/2008 | | Capsule | | | |
| Tablet | | | | | 250 mg | 2/1/2008 | |
| | 250 mg-125 mg | 2/1/2008 | | | 500 mg | 2/1/2008 | |
| | 500 mg-125 mg | 2/1/2008 | | Suspension Recon | | | |
| | 875 mg-125 mg | 2/1/2008 | | | 125 mg/5 mL | 2/1/2008 | |
| | | | | | 250 mg/5 mL | 2/1/2008 | |
| AMOXICILLIN | | | | ANAGRELIDE | | | |
| Oral | | | | Oral | | | |
| Capsule | | | | Capsule | | | |
| | 250 mg | 2/1/2008 | | | 0.5 mg | 5/1/2008 | |
| | 500 mg | 2/1/2008 | | | 1 mg | 5/1/2008 | |
| Suspension Recon | | | | ANASTROZOLE | | | |
| | 125 mg/5 mL | 2/1/2008 | | Oral | | | |
| | 200 mg/5 mL | 2/1/2008 | | Tablet | | | |
| | 250 mg/5 mL | 2/1/2008 | | | 1 mg | 10/1/2010 | |
| | 400 mg/5 mL | 2/1/2008 | | ANTIPYRINE and BENZOCAINE and GLYCERIN | | | |
| Tab Chew | | | | Otic | | | |
| | 125 mg | 2/1/2008 | | Drops | | | |
| | 200 mg | 2/1/2008 | | | 5.4 %-1.4 % | 2/1/2008 | |
| | 250 mg | 2/1/2008 | | ASPIRIN | | | |
| | 400 mg | 2/1/2008 | | Oral | | | |
| Tablet | | | | Tablet | | | |
| | 500 mg | 2/1/2008 | | | | | |
| | 875 mg | 2/1/2008 | | | | | |
| AMPHET ASP and AMPHET and D-AMPHET | | | | | | | |
| Oral | | | | | | | |
| Capsule (24hr) | | | | | | | |
| | 5 mg | 10/1/2010 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------------------|------------------------|----------|-----|---------------------------------|---------------------------|----------|-----|
| ASPIRIN | | | | AZITHROMYCIN | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Packet | | |
| | 325 mg | 2/1/2008 | | | 1 gram | 2/1/2008 | |
| | Tablet Chewable | | | | Suspension Recon | | |
| | 81 mg | 2/1/2008 | | | 100 mg/5 mL | 2/1/2008 | |
| | Tablet DR | | | | 200 mg/5 mL | 2/1/2008 | |
| | 81 mg | 2/1/2008 | | | Tablet | | |
| | 325 mg | 2/1/2008 | | | 250 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | 500 mg | 2/1/2008 | |
| | Tablet ER | | | | 600 mg | 2/1/2008 | |
| | 800 mg | 2/1/2008 | | | | | |
| | Rectal | | | | BACITRACIN | | |
| | Suppository | | | | Topical | | |
| | 600 mg | 1/1/2011 | | | Oinment | | |
| | | | | | 500 unit/gram | 2/1/2008 | |
| ATENOLOL | | | | BACITRACIN and POLYMYXIN | | | |
| | Oral | | | | Topical | | |
| | Tablet | | | | Oinment | | |
| | 25 mg | 2/1/2008 | | | 500 unit-10,000 unit/gram | 2/1/2008 | |
| | 50 mg | 2/1/2008 | | | | | |
| | 100 mg | 2/1/2008 | | | BACITRACIN ZINC | | |
| ATENOLOL and CHLORTHALIDONE | | | | | Topical | | |
| | Oral | | | | Oinment | | |
| | Tablet | | | | 500 unit/gram | 2/1/2008 | |
| | 50 mg-25 mg | 2/1/2008 | | | | | |
| | 100 mg-25 mg | 2/1/2008 | | | BACLOFEN | | |
| ATROPINE | | | | | Oral | | |
| | Ophthalmic | | | | Tablet | | |
| | Drops | | | | 10 mg | 2/1/2008 | |
| | 1 % | 2/1/2008 | | | 20 mg | 2/1/2008 | |
| | Oinment | | | | BALSALAZIDE | | |
| | 1 % | 2/1/2008 | | | Oral | | |
| | Oral | | | | Capsule | | |
| | Tablet | | | | 750 mg | 9/1/2009 | |
| | 0.4 mg | 2/1/2008 | | | BENZAEPRIIL | | |
| AZATHIOPRINE | | | | | Oral | | |
| | Oral | | | | Tablet | | |
| | Tablet | | | | 5 mg | 2/1/2008 | |
| | 50 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| AZITHROMYCIN | | | | | 20 mg | 2/1/2008 | |
| | Oral | | | | 40 mg | 2/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|---------------------------------------------------------|----------------|----------|-----|-------------------------------------------|-------------------|-----------|-----|
| BENAZEPRIL and HYDROCHLOROTHIAZIDE | | | | BETAMETHASONE VALERATE | | | |
| | Oral | | | | Topical | | |
| | Tablet | | | | Cream | | |
| | 5 mg-6.25 mg | 2/1/2008 | | | 0.1 % | 2/1/2008 | |
| | 10 mg-12.5 mg | 2/1/2008 | | | Lotion | | |
| | 20 mg-12.5 mg | 2/1/2008 | | | 0.1 % | 2/1/2008 | |
| | 20 mg-25 mg | 2/1/2008 | | | Oinment | | |
| | | | | | 0.1 % | 2/1/2008 | |
| BENZOCAINE | | | | BICALUTAMIDE | | | |
| | Otic | | | | Oral | | |
| | Drops | | | | Tablet | | |
| | 20 % | 2/1/2008 | | | 50 mg | 8/1/2009 | |
| BENZOYL PEROXIDE | | | | BISMUTH SUBSALICYLATE (Age 0-20) | | | |
| | Topical | | | | Oral | | |
| | Gel | | | | Suspension | | |
| | 10 % | 7/1/2009 | | | 262 mg/15 mL | 2/1/2008 | |
| | 5 % | 7/1/2009 | | | 525 mg/15 mL | 2/1/2008 | |
| BENZTROPINE | | | | BISOPROLOL | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 0.5 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| | 1 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | 2 mg | 2/1/2008 | | BISOPROLOL and HYDROCHLOROTHIAZIDE | | | |
| BETAMETHASONE DIPROPIONATE | | | | BISOPROLOL and HYDROCHLOROTHIAZIDE | | | |
| | Topical | | | | Oral | | |
| | Cream | | | | Tablet | | |
| | 0.05 % | 2/1/2008 | | | 10 mg-6.25 mg | 2/1/2008 | |
| | Gel | | | | 2.5 mg-6.25 mg | 2/1/2008 | |
| | 0.05 % | 2/1/2008 | | | 5 mg-6.25 mg | 2/1/2008 | |
| | Lotion | | | BRIMONIDINE | | | |
| | 0.05 % | 2/1/2008 | | | Ophthalmic | | |
| | Oinment | | | | Drops | | |
| | 0.05 % | 2/1/2008 | | | 0.15 % | 10/1/2010 | |
| BETAMETHASONE DIPROPIONATE and PROPOLYENE GLYCOL | | | | | 0.2 % | 2/1/2008 | |
| | Topical | | | BROMOCRIPTINE | | | |
| | Cream | | | | | | |
| | 0.05 % | 2/1/2008 | | | | | |
| | Oinment | | | | | | |
| | 0.05 % | 2/1/2008 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----------------------------|----------------------|-----------|-----|--------------------------------------------------|------------------------|-----------|-----|
| BROMOCRIPTINE | | | | CALAMINE COMBINATIONS (Age 0-20) | | | |
| | Oral | | | | Topical | | |
| | Capsule | | | | Lotion | 1/1/2011 | |
| | 5 mg | 7/1/2009 | | | Suspension | | |
| | Tablet | | | | 8 %-8 % | 1/1/2011 | |
| | 2.5 mg | 5/1/2008 | | CALCITONIN-SALMON | | | |
| BUDESONIDE (Age 0-8) | | | | | Nasal | | |
| | Inhalation | | | | Spray | 10/1/2009 | |
| | Ampul-Neb | | | | 200 unit/dose | | |
| | 0.25 mg/2 mL | 4/1/2010 | | CALCITRIOL | | | |
| | 0.5 mg/2 mL | 4/1/2010 | | | Oral | | |
| BUMETANIDE | | | | | Capsule | | |
| | Oral | | | | 0.25 mcg | 2/1/2008 | |
| | Tablet | | | | 0.5 mcg | 2/1/2008 | |
| | 0.5 mg | 2/1/2008 | | CALCIUM ACETATE | | | |
| | 1 mg | 2/1/2008 | | | Oral | | |
| | 2 mg | 2/1/2008 | | | Capsule | | |
| BUPROPION | | | | | 667 mg | 4/1/2009 | |
| | Oral | | | CALCIUM CARBONATE | | | |
| | Tablet | | | | Oral | | |
| | 75 mg | 2/1/2008 | | | Tablet Chewable | | |
| | 100 mg | 2/1/2008 | | | 320 mg (750 mg) | 1/1/2011 | |
| | Tablet (24hr) | | | | 400 mg | 1/1/2011 | |
| | 150 mg | 11/1/2010 | | | 400 mg (1,000 mg) | 1/1/2011 | |
| | 300 mg | 2/1/2008 | | | 600 mg (1.5 gram) | 1/1/2011 | |
| | Tablet ER | | | CALCIUM CARBONATE and MAGNESIUM HYDROXIDE | | | |
| | 100 mg | 2/1/2008 | | | Oral | | |
| | 150 mg | 5/1/2008 | | | Suspension | | |
| | 150 mg | 2/1/2008 | | | 400 mg-135 mg/5 mL | 1/1/2011 | |
| | 200 mg | 2/1/2008 | | | Tablet Chewable | | |
| BUSPIRONE | | | | | 700 mg-300 mg | 1/1/2011 | |
| | Oral | | | CAPSAICIN | | | |
| | Tablet | | | | Topical | | |
| | 5 mg | 2/1/2008 | | | Cream | | |
| | 7.5 mg | 2/1/2008 | | | 0.025 % | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | | 0.075 % | 2/1/2008 | |
| | 15 mg | 2/1/2008 | | | 0.1 % | 10/1/2010 | |
| | 30 mg | 2/1/2008 | | BUTORPHANOL | | | |
| | Nasal | | | | Nasal | | |
| | Spray | | | | Spray | | |
| | 10 mg/mL | 5/1/2008 | | | 10 mg/mL | 5/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------------------------|--------------|-----------|-----|-------------------------------|-------------|----------|-----|
| CAPTOPRIL | | | | CARBOXYMETHYLCELLULOSE | | | |
| Oral | | | | Ophthalmic | | | |
| Tablet | | | | Drops | | | |
| | 12.5 mg | 2/1/2008 | | | 0.5 % | 1/1/2011 | |
| | 25 mg | 2/1/2008 | | | 1 % | 1/1/2011 | |
| | 50 mg | 2/1/2008 | | CARTEOLOL | | | |
| | 100 mg | 2/1/2008 | | Ophthalmic | | | |
| CAPTOPRIL and HYDROCHLOROTHIAZIDE | | | | Drops | | | |
| Oral | | | | 1 % | | | |
| Tablet | | | | 2/1/2008 | | | |
| | 25 mg-15 mg | 2/1/2008 | | CARVEDILOL | | | |
| | 25 mg-25 mg | 2/1/2008 | | Oral | | | |
| | 50 mg-15 mg | 2/1/2008 | | Tablet | | | |
| | 50 mg-25 mg | 2/1/2008 | | | 3.125 mg | 2/1/2008 | |
| CARBAMAZEPINE | | | | | 6.25 mg | 2/1/2008 | |
| Oral | | | | | 12.5 mg | 2/1/2008 | |
| Suspension | | | | | 25 mg | 2/1/2008 | |
| | 100 mg/5 mL | 2/1/2008 | | CASTOR (Age 0-20) | | | |
| | 200 mg/10 mL | 2/1/2008 | | Oral | | | |
| Tablet | | | | Oil | | | |
| | 200 mg | 2/1/2008 | | | 100 % | 1/1/2011 | |
| Tablet Chewable | | | | | | 1/1/2011 | |
| | 100 mg | 2/1/2008 | | CEFACLOR | | | |
| CARBIDOPA and LEVODOPA | | | | Oral | | | |
| Oral | | | | Capsule | | | |
| Tablet | | | | | 250 mg | 2/1/2008 | |
| | 10 mg-100 mg | 2/1/2008 | | | 500 mg | 2/1/2008 | |
| | 25 mg-100 mg | 2/1/2008 | | Suspension Recon | | | |
| | 25 mg-250 mg | 2/1/2008 | | | 125 mg/5 mL | 2/1/2008 | |
| Tablet ER | | | | | 250 mg/5 mL | 2/1/2008 | |
| | 25 mg-100 mg | 2/1/2008 | | | 375 mg/5 mL | 2/1/2008 | |
| | 50 mg-200 mg | 2/1/2008 | | CEFADROXIL | | | |
| Tablet Rapdis | | | | Oral | | | |
| | 10 mg-100 mg | 11/1/2008 | | Capsule | | | |
| | 25 mg-100 mg | 11/1/2008 | | | 500 mg | 2/1/2008 | |
| | 25 mg-250 mg | 11/1/2008 | | Suspension Recon | | | |
| CARBOXYMETHYLCELLULOSE | | | | | 250 mg/5 mL | 2/1/2008 | |
| Ophthalmic | | | | | 500 mg/5 mL | 2/1/2008 | |
| Droperette | | | | Tablet | | | |
| | 0.5 % | 1/1/2011 | | | 1 gram | 2/1/2008 | |
| | 1 % | 1/1/2011 | | CEFDINIR | | | |
| Drops | | | | Oral | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------|-------------------------|-----------|-----|-------------------------|------------------------|----------|-----|
| CEFDINIR | | | | CHLORAL HYDRATE | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Syrup | | |
| | 300 mg | 9/1/2009 | | | 500 mg/5 mL | 2/1/2008 | |
| | Suspension Recon | | | | | | |
| | 250 mg/5 mL | 5/1/2008 | | | | | |
| CEFPROZIL | | | | CHLORDIAZEPOXIDE | | | |
| | Oral | | | | Oral | | |
| | Suspension Recon | | | | Capsule | | |
| | 125 mg/5 mL | 5/1/2008 | | | 5 mg | 2/1/2008 | |
| | 250 mg/5 mL | 5/1/2008 | | | 10 mg | 2/1/2008 | |
| | Tablet | | | | 25 mg | 2/1/2008 | |
| | 250 mg | 5/1/2008 | | | | | |
| | 500 mg | 5/1/2008 | | | | | |
| CEFUROXIME | | | | CHLORHEXIDINE | | | |
| | Oral | | | | Mucous Membrane | | |
| | Suspension Recon | | | | Mouthwash | | |
| | 125 mg/5 mL | 10/1/2010 | | | 0.12 % | 2/1/2008 | |
| | 250 mg/5 mL | 7/1/2009 | | | | | |
| | Tablet | | | | | | |
| | 250 mg | 2/1/2008 | | | | | |
| | 500 mg | 2/1/2008 | | | | | |
| CEPHALEXIN | | | | CHLOROTHIAZIDE | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Tablet | | |
| | 250 mg | 2/1/2008 | | | 250 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | 500 mg | 2/1/2008 | |
| | Suspension Recon | | | | | | |
| | 125 mg/5 mL | 2/1/2008 | | | | | |
| | 250 mg/5 mL | 2/1/2008 | | | | | |
| | Tablet | | | | | | |
| | 500 mg | 2/1/2008 | | | | | |
| CETIRIZINE | | | | CHLORPROMAZINE | | | |
| | Oral | | | | Oral | | |
| | Solution | | | | Tablet | | |
| | 1 mg/mL | 4/1/2010 | | | 10 mg | 2/1/2008 | |
| | Tablet | | | | 25 mg | 2/1/2008 | |
| | 5 mg | 3/1/2008 | | | 50 mg | 2/1/2008 | |
| | 10 mg | 3/1/2008 | | | 100 mg | 2/1/2008 | |
| | | | | | 200 mg | 2/1/2008 | |
| CHLORAL HYDRATE | | | | CHLORPROPAMIDE | | | |
| | Oral | | | | Oral | | |
| | | | | | Tablet | | |
| | | | | | 100 mg | 5/1/2008 | |
| | | | | | 250 mg | 2/1/2008 | |
| CHLORAL HYDRATE | | | | CHLORTHALIDONE | | | |
| | Oral | | | | Oral | | |
| | | | | | Tablet | | |
| | | | | | 15 mg | 2/1/2008 | |
| | | | | | 25 mg | 2/1/2008 | |
| | | | | | 50 mg | 2/1/2008 | |
| CHLORAL HYDRATE | | | | CHLORZOXAZONE | | | |
| | Oral | | | | Oral | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----|-----------------------------------------|----------|-----|-----|---------------------------------------|----------|-----|
| | CHLORZOAZONE | | | | CIMETIDINE | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Solution | | |
| | 500 mg | 2/1/2008 | | | 300 mg/5 mL | 2/1/2008 | |
| | CHOLECALCIFEROL (Age 0-20) | | | | Tablet | | |
| | Oral | | | | 200 mg | 2/1/2008 | |
| | Drops | | | | 300 mg | 2/1/2008 | |
| | 400 U/ml | 8/1/2011 | | | 400 mg | 2/1/2008 | |
| | Tablet | | | | 800 mg | 2/1/2008 | |
| | 1000 Units | 8/1/2011 | | | CIPROFLOXACIN | | |
| | CHOLESTYRAMINE and ASPARTAME | | | | Oral | | |
| | Oral | | | | Tablet | | |
| | Packet | | | | 100 mg | 2/1/2008 | |
| | 4 gram | 2/1/2008 | | | 250 mg | 2/1/2008 | |
| | Powder | | | | 500 mg | 2/1/2008 | |
| | 4 gram | 2/1/2008 | | | 750 mg | 2/1/2008 | |
| | CHOLESTYRAMINE and SUCROSE | | | | CITALOPRAM | | |
| | Oral | | | | Oral | | |
| | Packet | | | | Solution | | |
| | 4 gram | 2/1/2008 | | | 10 mg/5 mL | 2/1/2008 | |
| | Powder | | | | Tablet | | |
| | 4 gram | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | CHOLINE and MAGNESIUM SALICYLATE | | | | 20 mg | 2/1/2008 | |
| | Oral | | | | 40 mg | 2/1/2008 | |
| | Liquid | | | | CITRIC ACID and SODIUM CITRATE | | |
| | 500 mg/5 mL | 2/1/2008 | | | Oral | | |
| | Tablet | | | | Solution | | |
| | 500 mg | 2/1/2008 | | | 334 mg-500 mg/5 mL | 2/1/2008 | |
| | 750 mg | 2/1/2008 | | | CLARITHROMYCIN | | |
| | 1,000 mg | 2/1/2008 | | | Oral | | |
| | CILOSTAZOL | | | | Suspension Recon | | |
| | Oral | | | | 125 mg/5 mL | 5/1/2008 | |
| | Tablet | | | | 250 mg/5 mL | 5/1/2008 | |
| | 50 mg | 2/1/2008 | | | CLEMASTINE | | |
| | 100 mg | 2/1/2008 | | | Oral | | |
| | CIMETIDINE | | | | Syrup | | |
| | Oral | | | | 0.67 mg/5 mL | 2/1/2008 | |
| | Solution | | | | Tablet | | |
| | | | | | 2.68 mg | 2/1/2008 | |
| | CLINDAMYCIN | | | | Oral | | |
| | Oral | | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------------|------------|-----------|-----|----------------------------------------|----------|----------|-----|
| CLINDAMYCIN | | | | CLONAZEPAM | | | |
| Oral | | | | Oral | | | |
| Capsule | | | | Tablet Rapdis | | | |
| | 150 mg | 2/1/2008 | | | 0.125 mg | 2/1/2008 | |
| | 300 mg | 2/1/2008 | | | 0.25 mg | 2/1/2008 | |
| CLINDAMYCIN PALMITATE | | | | | | | |
| Oral | | | | | | | |
| Solution Recon | | | | | | | |
| | 75 mg/5 mL | 10/1/2010 | | | | | |
| CLINDAMYCIN PHOSPHATE | | | | CLONIDINE | | | |
| Topical | | | | Oral | | | |
| Gel | | | | Tablet | | | |
| | 1 % | 2/1/2008 | | | 0.1 mg | 2/1/2008 | |
| Lotion | | | | | | | |
| | 1 % | 2/1/2008 | | | 0.2 mg | 2/1/2008 | |
| Solution | | | | | | | |
| | 1 % | 2/1/2008 | | | 0.3 mg | 2/1/2008 | |
| CLOBETASOL | | | | CLORAZEPATE | | | |
| Topical | | | | Oral | | | |
| Cream | | | | Tablet | | | |
| | 0.05 % | 2/1/2008 | | | 3.75 mg | 5/1/2008 | |
| | 0.05 % | 2/1/2008 | | | 7.5 mg | 5/1/2008 | |
| Gel | | | | | | | |
| | 0.05 % | 2/1/2008 | | | 15 mg | 5/1/2008 | |
| Oinment | | | | | | | |
| | 0.05 % | 2/1/2008 | | | | | |
| Solution | | | | | | | |
| | 0.05 % | 2/1/2008 | | | | | |
| CLOMIPRAMINE | | | | CLOTTRIMAZOLE | | | |
| Oral | | | | Topical | | | |
| Capsule | | | | Cream | | | |
| | 25 mg | 2/1/2008 | | | 1 % | 2/1/2008 | |
| | 50 mg | 2/1/2008 | | Solution | | | |
| | 75 mg | 2/1/2008 | | | 1 % | 2/1/2008 | |
| CLONAZEPAM | | | | Vaginal | | | |
| Oral | | | | Cream | | | |
| Tablet | | | | | | | |
| | 0.5 mg | 2/1/2008 | | | 1 % | 2/1/2008 | |
| | 1 mg | 2/1/2008 | | | 2 % | 2/1/2008 | |
| | 2 mg | 2/1/2008 | | Tablet | | | |
| Tablet Rapdis | | | | | | | |
| | | | | | 100 mg | 2/1/2008 | |
| | | | | CLOTTRIMAZOLE and BETAMETHASONE | | | |
| | | | | Topical | | | |
| | | | | Cream | | | |
| | | | | | | | |
| | | | | 1 %-0.05 % | | | |
| | | | | 2/1/2008 | | | |
| | | | | Lotion | | | |
| | | | | | | | |
| | | | | 1 %-0.05 % | | | |
| | | | | 2/1/2008 | | | |
| | | | | CLOZAPINE | | | |
| | | | | Oral | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|---------------------------------|--------------------|----------|-----|-------------------------------|------------|----------|-----|
| CLOZAPINE | | | | CYCLOPENTOLATE | | | |
| | Oral | | | | Ophthalmic | | |
| | Tablet | | | | Drops | | |
| | 12.5 mg | 2/1/2008 | | | 1 % | 2/1/2008 | |
| | 25 mg | 2/1/2008 | | | | | |
| | 50 mg | 2/1/2008 | | | | | |
| | 100 mg | 2/1/2008 | | | | | |
| | 200 mg | 2/1/2008 | | | | | |
| COD LIVER OIL/ZINC OXIDE | | | | CYCLOPHOSPHAMIDE | | | |
| | Topical | | | | Oral | | |
| | Ointment | | | | Tablet | | |
| | 40% | 8/1/2011 | | | 25 mg | 7/1/2009 | |
| | | | | | 50 mg | 7/1/2009 | |
| CODEINE | | | | CYCLOSPORINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Capsule | | |
| | 15 mg | 2/1/2008 | | | 100 mg | 6/1/2009 | |
| | 30 mg | 2/1/2008 | | | 25 mg | 6/1/2009 | |
| | 60 mg | 2/1/2008 | | | Solution | | |
| | | | | | 100 mg/mL | 6/1/2009 | |
| CODEINE and PROMETHAZINE | | | | CYCLOSPORINE, MODIFIED | | | |
| | Oral | | | | Oral | | |
| | Syrup | | | | Capsule | | |
| | 10 mg-6.25 mg/5 mL | 5/1/2008 | | | 25 mg | 5/1/2008 | |
| | | | | | 50 mg | 7/1/2009 | |
| | | | | | 100 mg | 5/1/2008 | |
| | | | | | Solution | | |
| | | | | | 100 mg/mL | 6/1/2009 | |
| COLCHICINE | | | | CYPROHEPTADINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Syrup | | |
| | 0.6 mg | 2/1/2008 | | | 2 mg/5 mL | 2/1/2008 | |
| | | | | | Tablet | | |
| | | | | | 4 mg | 2/1/2008 | |
| CROMOLYN SODIUM | | | | DANTROLENE | | | |
| | Inhalation | | | | Oral | | |
| | Ampul-Neb | | | | Capsule | | |
| | 20 mg/2 mL | 2/1/2008 | | | 25 mg | 5/1/2008 | |
| | | | | | 50 mg | 5/1/2008 | |
| | | | | | 100 mg | 5/1/2008 | |
| | Ophthalmic | | | | | | |
| | Drops | | | | | | |
| | 4 % | 2/1/2008 | | | | | |
| CYCLOBENZAPRINE | | | | DESIPRAMINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 5 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | | 25 mg | 2/1/2008 | |
| CYCLOPENTOLATE | | | | | | | |
| | Ophthalmic | | | | | | |
| N/D; N=New, D=Delete | | | | | | | |

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------------------------|------------------------|----------|-----|------------------------------------------------------------------------------|-------------------------|----------|-----|
| DESIPRAMINE | | | | DEXAMETHASONE | | | |
| Oral | | | | Oral | | | |
| Tablet | | | | Elixir | | | |
| | 50 mg | 2/1/2008 | | | 0.5 mg/5 mL | 2/1/2008 | |
| | 75 mg | 2/1/2008 | | Solution | | | |
| | 100 mg | 2/1/2008 | | | 0.5 mg/5 mL | 2/1/2008 | |
| | 150 mg | 2/1/2008 | | Tablet | | | |
| DESMOPRESSIN | | | | | 0.5 mg | 2/1/2008 | |
| Nasal | | | | | 0.75 mg | 2/1/2008 | |
| Spray | | | | | 1 mg | 2/1/2008 | |
| | 10 mcg/spray (0.1 mL) | 5/1/2008 | | | 1.5 mg | 2/1/2008 | |
| Nasal (non-refrig.) | | | | | 2 mg | 2/1/2008 | |
| Spray/Pump | | | | | 4 mg | 2/1/2008 | |
| | 10 mcg/spray (0.1 mL) | 5/1/2008 | | | 6 mg | 2/1/2008 | |
| Oral | | | | DEXMETHYLPHENIDATE | | | |
| Tablet | | | | ORAL | | | |
| | 0.1 mg | 5/1/2008 | | Tablet | | | |
| | 0.2 mg | 5/1/2008 | | | 10 mg | 8/1/2011 | |
| DESOGESTREL and ETHINYL ESTRADIOL | | | | | 2.5 mg | 8/1/2011 | |
| Oral | | | | | 5 mg | 8/1/2011 | |
| Tablet | | | | DEXTROAMPHETAMINE | | | |
| | 0.1 mg-25 mcg | 2/1/2008 | | Oral | | | |
| | (7)/0.125 mg-25 mcg | | | Capsule ER | | | |
| | (7)/0.15 mg-25 mcg (7) | | | | 5 mg | 3/1/2011 | |
| | 0.15 mg-0.03 mg | 2/1/2008 | | | 10 mg | 3/1/2011 | |
| | | | | | 15 mg | 3/1/2011 | |
| DESONIDE | | | | DEXTROMETHORPHAN | | | |
| Topical | | | | Oral | | | |
| Cream | | | | Liquid | | | |
| | 0.05 % | 2/1/2008 | | | 15 mg/5 mL | 1/1/2011 | |
| Lotion | | | | Syrup | | | |
| | 0.05 % | 2/1/2008 | | | 5 mg/5 mL | 1/1/2011 | |
| Ointment | | | | | 7.5 mg/5 mL | 1/1/2011 | |
| | 0.05 % | 2/1/2008 | | | 10 mg/5 mL | 1/1/2011 | |
| | | | | | 15 mg/5 mL | 1/1/2011 | |
| DEXAMETHASONE | | | | DEXTROMETHORPHAN TAN and PSEUDOEPHENDRINE TAN and DEXCHLORPHENIRAMINE | | | |
| Ophthalmic | | | | Oral | | | |
| Drops | | | | Suspension | | | |
| | 0.1 % | 2/1/2008 | | | 25 mg-75 mg-2.5 mg/5 mL | 2/1/2008 | |
| Oral | | | | | | | |
| Drops | | | | | | | |
| | 0.5 mg/0.5 mL | 2/1/2008 | | | | | |
| Elixir | | | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------------------------|-----------------------------|----------|-----|------------------------------|-----------------------|----------|-----|
| DEXTROMETHROPHAN and PROMETHAZINE | | | | DICYCLOMINE | | | |
| | Oral | | | | Oral | | |
| | Syrup | | | | Syrup | | |
| | 15 mg-6.25 mg/5 mL | 2/1/2008 | | | 10 mg/5 mL | 2/1/2008 | |
| | | | | | Tablet | | |
| | | | | | 20 mg | 2/1/2008 | |
| DIAZEPAM | | | | DIDANOSINE | | | |
| | Oral | | | | Oral | | |
| | Concentrate | | | | Capsule DR | | |
| | 5 mg/mL | 2/1/2008 | | | 125 mg | 1/1/2011 | |
| | Solution | | | | 200 mg | 1/1/2011 | |
| | 5 mg/5 mL | 2/1/2008 | | | 250 mg | 8/2/2010 | |
| | Tablet | | | | 400 mg | 8/2/2010 | |
| | 2 mg | 2/1/2008 | | DIFLORASONE DIACETATE | | | |
| | 5 mg | 2/1/2008 | | | Topical | | |
| | 10 mg | 2/1/2008 | | | Cream | | |
| | | | | | 0.05 % | 2/1/2008 | |
| | Rectal | | | DIGOXIN | | | |
| | | | | | Oral | | |
| | 2.5 mg | 6/1/2011 | | | Solution | | |
| | 5 mg-7.5 mg-10 mg | 6/1/2011 | | | 50 mcg/mL | 2/1/2008 | |
| | 12.5 mg-15 mg-17.5 mg-20 mg | 6/1/2011 | | | Tablet | | |
| DICLOFENAC POTASSIUM | | | | | 125 mcg | 2/1/2008 | |
| | Oral | | | | 250 mcg | 2/1/2008 | |
| | Tablet | | | DILTIAZEM | | | |
| | 50 mg | 2/1/2008 | | | Oral | | |
| DICLOFENAC SODIUM | | | | | Capsule (12hr) | | |
| | Oral | | | | 60 mg | 2/1/2008 | |
| | Tablet (24hr) | | | | 90 mg | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | | 120 mg | 2/1/2008 | |
| | Tablet DR | | | | Capsule (24hr) | | |
| | 25 mg | 2/1/2008 | | | 120 mg | 2/1/2008 | |
| | 50 mg | 2/1/2008 | | | 180 mg | 2/1/2008 | |
| | 75 mg | 2/1/2008 | | | 240 mg | 2/1/2008 | |
| | | | | | 300 mg | 2/1/2008 | |
| DICLOXACILLIN | | | | | Capsule ER | | |
| | Oral | | | | 120 mg | 2/1/2008 | |
| | Capsule | | | | 180 mg | 2/1/2008 | |
| | 250 mg | 2/1/2008 | | | 240 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | 300 mg | 2/1/2008 | |
| DICYCLOMINE | | | | | Capsule ER DEG | | |
| | Oral | | | | 120 mg | 2/1/2008 | |
| | Capsule | | | | 180 mg | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | | 240 mg | 2/1/2008 | |
| | | | | | 300 mg | 2/1/2008 | |
| | | | | | 360 mg | 2/1/2008 | |
| | | | | | 420 mg | 2/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----------------------------------|----------------------|----------|-----|-----------------------------------|--------|-----------|-----|
| DILTIAZEM | | | | DISOPYRAMIDE | | | |
| Oral | | | | Oral | | | |
| Capsule ER DEG | | | | Capsule ER | | | |
| | 120 mg | 2/1/2008 | | | 150 mg | 2/1/2008 | |
| | 180 mg | 2/1/2008 | | DISULFIRAM | | | |
| | 240 mg | 2/1/2008 | | Oral | | | |
| Tablet | | | | Tablet | | | |
| | 30 mg | 2/1/2008 | | | 250 mg | 7/1/2011 | |
| | 60 mg | 2/1/2008 | | DIVALPROEX | | | |
| | 90 mg | 2/1/2008 | | Oral | | | |
| | 120 mg | 2/1/2008 | | Capsule Sprinkle | | | |
| DIPHENHYDRAMINE | | | | Tablet (24hr) | | | |
| Oral | | | | | 125 mg | 10/1/2009 | |
| Capsule | | | | | 250 mg | 9/1/2009 | |
| | 25 mg | 2/1/2008 | | | 500 mg | 9/1/2009 | |
| | 50 mg | 2/1/2008 | | Tablet DR | | | |
| Liquid | | | | | 125 mg | 9/1/2008 | |
| | 12.5 mg/5 mL | 2/1/2008 | | | 250 mg | 9/1/2008 | |
| Syrup | | | | | 500 mg | 9/1/2008 | |
| | 12.5 mg/5 mL | 2/1/2008 | | DOCUSATE SODIUM (Age 0-20) | | | |
| DIPHENOXYLATE and ATROPINE | | | | Oral | | | |
| Oral | | | | Capsule | | | |
| Liquid | | | | | 50 mg | 1/1/2011 | |
| | 2.5 mg-0.025 mg/5 mL | 2/1/2008 | | Tablet | | | |
| Tablet | | | | | 100 mg | 1/1/2011 | |
| | 2.5 mg-0.025 mg | 2/1/2008 | | DONEPEZIL | | | |
| DIPIVEFRIN | | | | Oral | | | |
| Ophthalmic | | | | Tablet | | | |
| Drops | | | | | 5 mg | 2/1/2011 | |
| | 0.1 % | 2/1/2008 | | | 10 mg | 2/1/2011 | |
| DIPYRIDAMOLE | | | | DORZOLAMIDE | | | |
| Oral | | | | Ophthalmic | | | |
| Tablet | | | | Drops | | | |
| | 25 mg | 2/1/2008 | | | 2 % | 2/1/2009 | |
| | 50 mg | 2/1/2008 | | DOXAZOSIN | | | |
| | 75 mg | 2/1/2008 | | Oral | | | |
| DISOPYRAMIDE | | | | Tablet | | | |
| Oral | | | | | 1 mg | 2/1/2008 | |
| Capsule | | | | | 2 mg | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | | 4 mg | 2/1/2008 | |
| | 150 mg | 2/1/2008 | | | 8 mg | 2/1/2008 | |
| Capsule ER | | | | | | | |
| N/D; N=New, D=Delete | | | | | | | |

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------------------------|-------------------|-----------|-----|----------------------------------------------------------|--------------------------|-----------|-----|
| DOXEPIN | | | | ENOXAPARIN | | | |
| Oral | | | | Sub-Q | | | |
| Capsule | | | | Syringe | | | |
| | 10 mg | 2/1/2008 | | | 80 mg/0.8 mL | 10/1/2010 | |
| | 25 mg | 2/1/2008 | | | 100 mg/mL | 10/1/2010 | |
| | 50 mg | 2/1/2008 | | | 120 mg/0.8 mL | 10/1/2010 | |
| | 75 mg | 2/1/2008 | | | 150 mg/mL | 10/1/2010 | |
| | 100 mg | 2/1/2008 | | EPINEPHRINE (If generic available in marketplace) | | | |
| | 150 mg | 2/1/2008 | | Intramuscular | | | |
| Concentrate | | | | Pen Injector | | | |
| | 10 mg/mL | 2/1/2008 | | | 0.15 mg/0.15 mL (1:1000) | 10/1/2010 | |
| DOXYCYCLINE | | | | | 0.3 mg/0.3 mL (1:1,000) | 10/1/2010 | |
| Oral | | | | ERYTHROMYCIN | | | |
| Capsule | | | | Ophthalmic | | | |
| | 50 mg | 2/1/2008 | | Ointment | | | |
| | 100 mg | 2/1/2008 | | | 5 mg/gram (0.5 %) | 2/1/2008 | |
| Tablet | | | | Oral | | | |
| | 100 mg | 2/1/2008 | | Capsule DR | | | |
| DROSPIRENONE and ETHINYL ESTRADIOL | | | | | 250 mg | 2/1/2008 | |
| Oral | | | | Tablet | | | |
| | 0.02 mg-3 mg (24) | 10/1/2010 | | | 250 mg | 2/1/2008 | |
| ENALAPRIL | | | | | 500 mg | 2/1/2008 | |
| Oral | | | | Tablet DR | | | |
| Tablet | | | | | 250 mg | 2/1/2008 | |
| | 2.5 mg | 2/1/2008 | | | 333 mg | 2/1/2008 | |
| | 5 mg | 2/1/2008 | | | 500 mg | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | Topical | | | |
| | 20 mg | 2/1/2008 | | Gel | | | |
| ENALAPRIL and HYDROCHLOROTHIAZIDE | | | | | 2 % | 5/1/2008 | |
| Oral | | | | ERYTHROMYCIN and SULFISOXAZOLE | | | |
| Tablet | | | | Oral | | | |
| | 10 mg-25 mg | 2/1/2008 | | Suspension Recon | | | |
| | 5 mg-12.5 mg | 2/1/2008 | | | 200 mg-600 mg/5 mL | 2/1/2008 | |
| ENOXAPARIN | | | | ERYTHROMYCIN ETHYLSUCCINATE | | | |
| Sub-Q | | | | Oral | | | |
| Syringe | | | | Suspension Recon | | | |
| | 30 mg/0.3 mL | 10/1/2010 | | | 200 mg/5 mL | 2/1/2008 | |
| | 40 mg/0.4 mL | 10/1/2010 | | | 400 mg/5 mL | 2/1/2008 | |
| | 60 mg/0.6 mL | 10/1/2010 | | Tablet | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------------|-------------------------|----------|-----|---------------------|--------------------|----------|-----|
| FLUOCINOLONE | | | | FLUOXETINE | | | |
| | Topical | | | | Oral | | |
| | Oinment | | | | Tablet | | |
| | 0.025 % | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | Shampoo | | | | 20 mg | 2/1/2008 | |
| | 0.01 % | 2/1/2008 | | FLUPHENAZINE | | | |
| | Solution | | | | Oral | | |
| | 0.01 % | 2/1/2008 | | | Concentrate | | |
| FLUOCINONIDE | | | | | 5 mg/mL | 2/1/2008 | |
| | Topical | | | | Elixir | | |
| | Cream | | | | 2.5 mg/5 mL | 2/1/2008 | |
| | 0.05 % | 2/1/2008 | | | Tablet | | |
| | Gel | | | | 1 mg | 2/1/2008 | |
| | 0.05 % | 2/1/2008 | | | 2.5 mg | 2/1/2008 | |
| | Oinment | | | | 5 mg | 2/1/2008 | |
| | 0.05 % | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | Solution | | | FLURBIPROFEN | | | |
| | 0.05 % | 2/1/2008 | | | Oral | | |
| FLUOCINONIDE EMOLLIENT | | | | | Tablet | | |
| | Topical | | | | 50 mg | 2/1/2008 | |
| | Cream | | | | 100 mg | 2/1/2008 | |
| | 0.05 % | 2/1/2008 | | FLUTAMIDE | | | |
| FLUOROMETHOLONE | | | | | Oral | | |
| | Ophthalmic | | | | Capsule | | |
| | Drops Suspension | | | | 125 mg | 7/1/2009 | |
| | 0.1 % | 2/1/2008 | | FLUTICASONE | | | |
| FLUOROURACIL | | | | | Nasal | | |
| | Topical | | | | Spray Susp | | |
| | Cream | | | | 50 mcg | 2/1/2008 | |
| | 5 % | 5/1/2009 | | | Topical | | |
| | Solution | | | | Cream | | |
| | 2 % | 5/1/2009 | | | 0.05 % | 2/1/2008 | |
| | 5 % | 5/1/2009 | | | Oinment | | |
| FLUOXETINE | | | | | 0.005 % | 2/1/2008 | |
| | Oral | | | FLUVOXAMINE | | | |
| | Capsule | | | | Oral | | |
| | 10 mg | 2/1/2008 | | | Tablet | | |
| | 20 mg | 2/1/2008 | | | 25 mg | 2/1/2008 | |
| | 40 mg | 2/1/2008 | | | 50 mg | 2/1/2008 | |
| | Solution | | | | 100 mg | 2/1/2008 | |
| | 20 mg/5 mL | 2/1/2008 | | FOLIC ACID | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|--------------------|-------------|-----------|-----|----------------------------------|----------------|----------|-----|
| FOLIC ACID | | | | GENTAMICIN | | | |
| | Oral | | | | Topical | | |
| | Tablet | | | | Cream | | |
| | 1 mg | 2/1/2008 | | | 0.1 % | 2/1/2008 | |
| FUROSEMIDE | | | | | Ointment | | |
| | Oral | | | | 0.1 % | 2/1/2008 | |
| | Solution | | | GENTIAN VIOLET (Age 0-20) | | | |
| | 10 mg/mL | 2/1/2008 | | | Topical | | |
| | 40 mg/5 mL | 2/1/2008 | | | Solution | | |
| | Tablet | | | | 1 % | 1/1/2011 | |
| | 20 mg | 2/1/2008 | | GLIMEPIRIDE | | | |
| | 40 mg | 2/1/2008 | | | Oral | | |
| | 80 mg | 2/1/2008 | | | Tablet | | |
| GABAPENTIN | | | | | 1 mg | 2/1/2008 | |
| | Oral | | | | 2 mg | 2/1/2008 | |
| | Capsule | | | | 4 mg | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | GLIPIZIDE | | | |
| | 300 mg | 2/1/2008 | | | Oral | | |
| | 400 mg | 2/1/2008 | | | Tablet | | |
| | Solution | | | | 5 mg | 2/1/2008 | |
| N | 250 mg/5 mL | 10/1/2011 | | | 10 mg | 2/1/2008 | |
| | Tablet | | | | Tablet (24hr) | | |
| | 600 mg | 2/1/2008 | | | 2.5 mg | 2/1/2008 | |
| | 800 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| | | | | | 10 mg | 2/1/2008 | |
| GALANTAMINE | | | | GLIPIZIDE and METFORMIN | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 4 mg | 12/1/2008 | | | 2.5 mg-250 mg | 2/1/2008 | |
| | 8 mg | 12/1/2008 | | | 2.5 mg-500 mg | 2/1/2008 | |
| | 12 mg | 12/1/2008 | | | 5 mg-500 mg | 2/1/2008 | |
| GEMFIBROZIL | | | | GLYBURIDE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 600 mg | 2/1/2008 | | | 1.25 mg | 2/1/2008 | |
| GENTAMICIN | | | | | 2.5 mg | 2/1/2008 | |
| | Ophthalmic | | | | 5 mg | 2/1/2008 | |
| | Drops | | | GLYBURIDE and METFORMIN | | | |
| | 0.3 % | 2/1/2008 | | | Oral | | |
| | Ointment | | | | Tablet | | |
| | 0.3 % | 2/1/2008 | | | 1.25 mg-250 mg | 2/1/2008 | |
| | Topical | | | GLYBURIDE and METFORMIN | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----------------------------------------|-------------------|----------|-----|------------------------------------------------------------|------------------------|----------|-----|
| GLYBURIDE and METFORMIN | | | | GUAIFENESIN and DEXTROMETHORPHAN | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet (12hr) | | |
| | 2.5 mg-500 mg | 2/1/2008 | | | 1,200 mg-60 mg | 2/1/2008 | |
| | 5 mg-500 mg | 2/1/2008 | | | | | |
| GLYBURIDE, MICRONIZED | | | | GUAIFENESIN and DEXTROMETHORPHAN and PSEUDOEPHEDRIN | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet (12hr) | | |
| | 1.5 mg | 2/1/2008 | | | 800 mg-60 mg-90 mg | 2/1/2008 | |
| | 3 mg | 2/1/2008 | | | | | |
| | 6 mg | 2/1/2008 | | | | | |
| GRISEOFULVIN, MICRONIZED | | | | GUAIFENESIN and PHENYLEPHRINE | | | |
| | Oral | | | | Oral | | |
| | Suspension | | | | Tablet | | |
| | 125 mg/5 mL | 4/2/2008 | | | 200 mg-25 mg | 2/1/2008 | |
| GUAIFENESIN | | | | GUAIFENESIN and PHENYLEPHRINE and PYRIL | | | |
| | Oral | | | | Oral | | |
| | Liquid | | | | Suspension | | |
| | 50 mg/5 mL | 1/1/2011 | | | 100 mg-5 mg-30 mg/5 mL | 2/1/2008 | |
| | 100 mg/5 mL | 2/1/2008 | | | Tablet | | |
| | 200 mg/5 mL | 1/1/2011 | | | 200 mg-25 mg-60 mg | 2/1/2008 | |
| | Syrup | | | | | | |
| | 100 mg/5 mL | 2/1/2008 | | | | | |
| GUAIFENESIN and CODEINE | | | | GUAIFENESIN and PSEUDOEPHEDRINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Capsule (12hr) | | |
| | 300 mg-10 mg | 2/1/2008 | | | 250 mg-120 mg | 2/1/2008 | |
| GUAIFENESIN and DEXTROMETHORPHAN | | | | GUANFACINE | | | |
| | Oral | | | | Oral | | |
| | Elixir | | | | Tablet | | |
| | 200 mg-20 mg/5 mL | 2/1/2008 | | | 1 mg | 2/1/2008 | |
| | Liquid | | | | 2 mg | 2/1/2008 | |
| | 100 mg-10 mg/5 mL | 2/1/2008 | | | | | |
| | 100 mg-5 mg/5 mL | 2/1/2008 | | | | | |
| | 200 mg-10 mg/5 mL | 1/1/2011 | | | | | |
| | 300 mg-20 mg/5 mL | 1/1/2011 | | | | | |
| | Solution | | | | | | |
| | 225 mg-25 mg/5 mL | 2/1/2008 | | | | | |
| | Syrup | | | | | | |
| | 100 mg-10 mg/5 mL | 2/1/2008 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|--------------------------------------|-----------------------------|----------|-----|--------------------------------------|---------------|----------|-----|
| HALOBETASOL | | | | HYDROCODONE and ACETAMINOPHEN | | | |
| Topical | | | | Oral | | | |
| Cream | | | | Tablet | | | |
| | 0.05 % | 5/1/2008 | | | 2.5 mg-500 mg | 2/1/2008 | |
| Oinment | | | | | 5 mg-325 mg | 2/1/2008 | |
| | 0.05 % | 5/1/2008 | | | 5 mg-500 mg | 2/1/2008 | |
| HALOPERIDOL | | | | | 7.5 mg-325 mg | 2/1/2008 | |
| Oral | | | | | 7.5 mg-500 mg | 2/1/2008 | |
| Concentrate | | | | | 7.5 mg-650 mg | 2/1/2008 | |
| | 2 mg/mL | 2/1/2008 | | | 7.5 mg-750 mg | 2/1/2008 | |
| Tablet | | | | | 10 mg-325 mg | 2/1/2008 | |
| | 0.5 mg | 2/1/2008 | | | 10 mg-500 mg | 2/1/2008 | |
| | 1 mg | 2/1/2008 | | | 10 mg-650 mg | 2/1/2008 | |
| | 2 mg | 2/1/2008 | | | 10 mg-660 mg | 2/1/2008 | |
| | 5 mg | 2/1/2008 | | | 10 mg-750 mg | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | HYDROCORTISONE | | | |
| | 20 mg | 2/1/2008 | | Oral | | | |
| HYDRALAZINE | | | | Tablet | | | |
| Oral | | | | | 5 mg | 7/1/2009 | |
| Tablet | | | | | 10 mg | 7/1/2009 | |
| | 10 mg | 2/1/2008 | | | 20 mg | 2/1/2008 | |
| | 25 mg | 2/1/2008 | | Topical | | | |
| | 50 mg | 2/1/2008 | | Cream | | | |
| | 100 mg | 2/1/2008 | | | 0.5 % | 2/1/2008 | |
| HYDROCHLOROTHIAZIDE | | | | | 1 % | 2/1/2008 | |
| Oral | | | | | 2.5 % | 2/1/2008 | |
| Capsule | | | | Lotion | | | |
| | 12.5 mg | 2/1/2008 | | | 1 % | 2/1/2008 | |
| Tablet | | | | | 2.5 % | 2/1/2008 | |
| | 25 mg | 2/1/2008 | | Oinment | | | |
| | 50 mg | 2/1/2008 | | | 0.5 % | 2/1/2008 | |
| HYDROCODONE and ACETAMINOPHEN | | | | | 1 % | 2/1/2008 | |
| Oral | | | | | 2.5 % | 2/1/2008 | |
| Capsule | | | | HYDROCORTISONE ACETATE | | | |
| | 5 mg-500 mg | 2/1/2008 | | Topical | | | |
| Solution | | | | Cream | | | |
| | 2.5 mg-167 mg/5 mL | 2/1/2008 | | | 0.5 % | 2/1/2008 | |
| | 5 mg-334 mg/10 mL | 2/1/2008 | | | 1 % | 2/1/2008 | |
| | 7.5 mg-500 mg/15 mL | 2/1/2008 | | Oinment | | | |
| | 7.5 mg-500 mg/15 mL (15 mL) | 2/1/2008 | | | 1 % | 2/1/2008 | |
| Tablet | | | | HYDROCORTISONE VALERATE | | | |
| | | | | Topical | | | |
| | | | | Cream | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|--------------------------------|----------------------|----------|-----|-----------------------------|------------------------|-----------|-----|
| HYDROCORTISONE VALERATE | | | | HYOSCYAMINE | | | |
| | Topical | | | | Sublingual | | |
| | Cream | | | | Tablet | | |
| | 0.2 % | 2/1/2008 | | | 0.125 mg | 2/1/2008 | |
| HYDROMORPHONE | | | | HYPROMELLOSE | | | |
| | Oral | | | | Ophthalmic | | |
| | Tablet | | | | Drops | | |
| | 2 mg | 5/1/2008 | | | 0.3 % | 1/1/2011 | |
| | 4 mg | 5/1/2008 | | | 0.4 % | 1/1/2011 | |
| | 8 mg | 5/1/2008 | | | Gel | | |
| HYDROXYCHLOROQUINE | | | | | 0.3 % | 1/1/2011 | |
| | Oral | | | IBUPROFEN | | | |
| | Tablet | | | | Oral | | |
| | 200 mg | 2/1/2008 | | | Suspension | | |
| HYDROXYUREA | | | | | 100 mg/5 mL | 2/1/2008 | |
| | Oral | | | | Tablet | | |
| | Capsule | | | | 200 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | 400 mg | 2/1/2008 | |
| HYDROXYZINE | | | | | 600 mg | 2/1/2008 | |
| | Oral | | | | 800 mg | 2/1/2008 | |
| | Capsule | | | IBUPROFEN (Age 0-12) | | | |
| | 25 mg | 2/1/2008 | | | Oral | | |
| | 50 mg | 2/1/2008 | | | Tablet Chewable | | |
| | 100 mg | 2/1/2008 | | | 100 mg | 2/1/2008 | |
| | Syrup | | | IMIPRAMINE | | | |
| | 10 mg/5 mL | 2/1/2008 | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 10 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | 25 mg | 2/1/2008 | | | 25 mg | 2/1/2008 | |
| | 50 mg | 2/1/2008 | | | 50 mg | 2/1/2008 | |
| HYOSCYAMINE | | | | IMIQUIMOD | | | |
| | Oral | | | | Topical | | |
| | Drops | | | | Cream | | |
| | 0.125 mg/mL | 2/1/2008 | | | 5 % | 10/1/2010 | |
| | Elixir | | | INDAPAMIDE | | | |
| | 125 mcg/5 mL | 2/1/2008 | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 0.125 mg | 2/1/2008 | | | 1.25 mg | 2/1/2008 | |
| | Tablet Rapdis | | | | 2.5 mg | 2/1/2008 | |
| | 0.125 mg | 2/1/2008 | | INDOMETHACIN | | | |
| | Sublingual | | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|----------------------------------|--------------------------------|-----------|-----|-------------------------------|----------------------|----------|-----|
| INDOMETHACIN | | | | ISONIAZID | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Tablet | | |
| | 25 mg | 2/1/2008 | | | 100 mg | 2/1/2008 | |
| | 50 mg | 2/1/2008 | | | 300 mg | 2/1/2008 | |
| | Capsule ER | | | ISOSORBIDE DINITRATE | | | |
| | 75 mg | 2/1/2008 | | | Oral | | |
| INSULIN - HUMULIN 70/30 | | | | | Tablet | | |
| | Sub-Q | | | | 5 mg | 2/1/2008 | |
| | Insulin Pen | | | | 10 mg | 2/1/2008 | |
| | 100 unit/mL (70-30) | 2/1/2008 | | | 20 mg | 2/1/2008 | |
| | Vial | | | | 30 mg | 2/1/2008 | |
| | 100 unit/mL (70-30) | 2/1/2008 | | | Tablet ER | | |
| INSULIN - HUMULIN N | | | | | 40 mg | 2/1/2008 | |
| | Sub-Q | | | | Sublingual | | |
| | Insulin Pen | | | | Sublingual | | |
| | 100 unit/mL (3 mL) | 2/1/2008 | | | 2.5 mg | 2/1/2008 | |
| | Vial | | | | 5 mg | 2/1/2008 | |
| | 100 unit/mL | 2/1/2008 | | ISOSORBIDE MONONITRATE | | | |
| INSULIN - HUMULIN R | | | | | Oral | | |
| | Injection | | | | Tablet | | |
| | Vial | | | | 10 mg | 2/1/2008 | |
| | 100 unit/mL | 2/1/2008 | | | 20 mg | 2/1/2008 | |
| IPRATROPIUM | | | | | Tablet (24hr) | | |
| | Inhalation | | | | 30 mg | 2/1/2008 | |
| | Solution | | | | 60 mg | 2/1/2008 | |
| | 0.2 mg/mL (0.02 %) | 2/1/2008 | | | 120 mg | 2/1/2008 | |
| | Nasal | | | ISOTRETINOIN | | | |
| | Spray | | | | Oral | | |
| | 21 mcg | 2/1/2008 | | | Capsule | | |
| | 42 mcg | 2/1/2008 | | | 10 mg | 8/2/2010 | |
| IPRATROPIUM and ALBUTEROL | | | | | 20 mg | 8/2/2010 | |
| | Inhalation | | | | 40 mg | 8/2/2010 | |
| | Ampul-Neb | | | KETOCONAZOLE | | | |
| | 0.5 mg-3 mg (2.5 mg base)/3 mL | 10/1/2010 | | | Oral | | |
| ISONIAZID | | | | | Tablet | | |
| | Oral | | | | 200 mg | 2/1/2008 | |
| | Syrup | | | | Topical | | |
| | 50 mg/5 mL | 2/1/2008 | | | Cream | | |
| | | | | | 2 % | 7/1/2011 | |
| | | | | | Shampoo | | |
| | | | | | 2 % | 2/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------------|-------------------|----------|-----|-----------------------------------|---------------------|-----------|-----|
| KETOPROFEN | | | | LEFLUNOMIDE | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Tablet | | |
| | 50 mg | 2/1/2008 | | | 10 mg | 2/1/2011 | |
| | 75 mg | 2/1/2008 | | | 20 mg | 2/1/2011 | |
| KETOROLAC TROMETHAMINE | | | | LETROZOLE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 10 mg | 2/1/2008 | | | 2.5 mg | 8/1/2011 | |
| LABETALOL | | | | LEVETIRACETAM | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Solution | | |
| | 100 mg | 2/1/2008 | | | 100 mg/mL | 2/1/2009 | |
| | 200 mg | 2/1/2008 | | | Tablet | | |
| | 300 mg | 2/1/2008 | | | 250 mg | 1/1/2009 | |
| LACTASE (Age 0-20) | | | | | 500 mg | 12/1/2008 | |
| | Oral | | | | 750 mg | 12/1/2008 | |
| | Capsule | | | | 1,000 mg | 2/1/2009 | |
| | 250 mg | 1/1/2011 | | LEVOBUNOLOL | | | |
| | Tablet | | | | Ophthalmic | | |
| | 3,000 unit | 1/1/2011 | | | Drops | | |
| LACTULOSE | | | | | 0.25 % | 2/1/2008 | |
| | Oral | | | | 0.5 % | 2/1/2008 | |
| | Solution | | | LEVOCARNITINE | | | |
| | 10 gram/15 mL | 2/1/2008 | | | Oral | | |
| | 10 gram/15 mL | 2/1/2008 | | | Solution | | |
| LAMOTRIGINE | | | | | 100 mg/mL | 1/1/2009 | |
| | Oral | | | LEVOFLOXACIN | | | |
| | Tab Disper | | | | Oral | | |
| | 25 mg | 4/1/2009 | | | Tablet | | |
| | 5 mg | 4/1/2009 | | | 250 mg | 8/1/2011 | |
| | Tablet | | | | 500 mg | 8/1/2011 | |
| | 25 mg | 4/1/2009 | | | 750 mg | 8/1/2011 | |
| | 100 mg | 4/1/2009 | | LEVONORGESTREL and ETHINYL | | | |
| | 150 mg | 4/1/2009 | | ESTRADIOL | | | |
| | 200 mg | 4/1/2009 | | | Oral | | |
| LATANOPROST | | | | | Tablet | | |
| | Ophthalmic | | | | 0.05 mg-0.03mg and | 2/1/2008 | |
| | Drops | | | | 0.075 mg-0.04mg and | | |
| | 0.005 % | 5/1/2011 | | | 0.125 mg-0.03mg | | |
| | | | | N | 0.09 mg-0.02 mg | 10/1/2011 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|---------------------------------------------|-------------------------------------|-----------|-----|-------------------------------------------|---------------|----------|-----|
| LEVONORGESTREL and ETHINYL ESTRADIOL | | | | LISINOPRIL | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 0.1 mg-0.02 mg | 2/1/2008 | | | 20 mg | 2/1/2008 | |
| | 0.15 mg-0.03 mg | 2/1/2008 | | | 30 mg | 2/1/2008 | |
| N | 0.15 mg-0.03 mg and 0.00 mg-0.01 mg | 10/1/2011 | | | 40 mg | 2/1/2008 | |
| | Tablet Dose Pak (3m) | | | LISINOPRIL and HYDROCHLOROTHIAZIDE | | | |
| | 0.15 mg-0.03 mg | 2/1/2008 | | | Oral | | |
| LEVOTHYROXINE | | | | | Tablet | | |
| | Oral | | | | 10 mg-12.5 mg | 2/1/2008 | |
| | Tablet | | | | 20 mg-12.5 mg | 2/1/2008 | |
| | 25 mcg | 2/1/2008 | | | 20 mg-25 mg | 2/1/2008 | |
| | 50 mcg | 2/1/2008 | | LITHIUM | | | |
| | 75 mcg | 2/1/2008 | | | Oral | | |
| | 88 mcg | 2/1/2008 | | | Capsule | | |
| | 100 mcg | 2/1/2008 | | | 150 mg | 2/1/2008 | |
| | 112 mcg | 2/1/2008 | | | 300 mg | 2/1/2008 | |
| | 125 mcg | 2/1/2008 | | | 600 mg | 2/1/2008 | |
| | 137 mcg | 2/1/2008 | | | Solution | | |
| | 150 mcg | 2/1/2008 | | | 8 mEq/5 mL | 2/1/2008 | |
| | 175 mcg | 2/1/2008 | | | Tablet | | |
| | 200 mcg | 2/1/2008 | | | 300 mg | 2/1/2008 | |
| | 300 mcg | 2/1/2008 | | | Tablet ER | | |
| LIDOCAINE | | | | | 300 mg | 2/1/2008 | |
| | Mucous Membrane | | | | 450 mg | 2/1/2008 | |
| | Jel | | | LOPERAMIDE | | | |
| | 2 % | 2/1/2008 | | | Oral | | |
| | 2 % | 2/1/2008 | | | Capsule | | |
| | Solution | | | | 2 mg | 2/1/2008 | |
| | 20 mg/mL | 2/1/2008 | | LOPERAMIDE (Age 0-20) | | | |
| LIPASE and PROTEASE and AMYLASE | | | | | Oral | | |
| | Oral | | | | Liquid | | |
| | Capsule DR | | | | 1 mg/5 mL | 2/1/2008 | |
| | 5,000 unit-17,000 unit-27,000 unit | 7/1/2010 | | | Tablet | | |
| LISINOPRIL | | | | | 2 mg | 2/1/2008 | |
| | Oral | | | LORATADINE | | | |
| | Tablet | | | | Oral | | |
| | 2.5 mg | 2/1/2008 | | | Solution | | |
| | 5 mg | 2/1/2008 | | | 5 mg/5 mL | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | | Tablet | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----------------------------------------|----------------|----------|-----|---------------------------------------------------|----------------------------|----------|-----|
| LORATADINE | | | | LOXAPINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Capsule | | |
| | 10 mg | 2/1/2008 | | | 25 mg | 5/1/2008 | |
| | | | | | 50 mg | 5/1/2008 | |
| LORATADINE and PSEUDOEPHEDRINE | | | | MAGALDRATE | | | |
| | Oral | | | | Oral | | |
| | Tablet (12hr) | | | | Suspension | | |
| | 5 mg-120 mg | 2/1/2008 | | | 540 mg/5 mL | 2/1/2008 | |
| | Tablet (24hr) | | | | | | |
| | 10 mg-240 mg | 2/1/2008 | | MAGNESIUM (Age 0-20) | | | |
| LORAZEPAM | | | | | Oral | | |
| | Oral | | | | Tablet | | |
| | Tablet | | | | 250 mg | 1/1/2011 | |
| | 0.5 mg | 2/1/2008 | | MAGNESIUM and ALUMINUM HYDROXIDE | | | |
| | 1 mg | 2/1/2008 | | | Oral | | |
| | 2 mg | 2/1/2008 | | | Suspension | | |
| LOSARTAN | | | | | 200 mg-200 mg/5 mL | 1/1/2011 | |
| | Oral | | | | 200 mg-225 mg/5 mL | 2/1/2008 | |
| | Tablet | | | | 500 mg-500 mg/5 mL | 1/1/2011 | |
| | 100 mg | 6/1/2010 | | | Tablet Chewable | | |
| | 25 mg | 6/1/2010 | | | 150 mg-300 mg | 1/1/2011 | |
| | 50 mg | 6/1/2010 | | MAGNESIUM CARBONATE and ALUMINUM HYDROXIDE | | | |
| LOSARTAN and HYDROCHLOROTHIAZIDE | | | | | Oral | | |
| | Oral | | | | Tablet Chewable | | |
| | Tablet | | | | 105 mg-160 mg | 1/1/2011 | |
| | 100 mg-12.5 mg | 6/1/2010 | | MAGNESIUM CHLORIDE (Age 0-20) | | | |
| | 100 mg-25 mg | 6/1/2010 | | | Oral | | |
| | 50 mg-12.5 mg | 6/1/2010 | | | Tablet ER | | |
| LOVASTATIN | | | | | 64 mg | 1/1/2011 | |
| | Oral | | | MAGNESIUM GLUCONATE (Age 0-20) | | | |
| | Tablet | | | | Oral | | |
| | 10 mg | 2/1/2008 | | | Liquid | | |
| | 20 mg | 2/1/2008 | | | 54 mg/5 ml (1,000 mg/5 mL) | 1/1/2011 | |
| | 40 mg | 2/1/2008 | | | Tablet | | |
| LOXAPINE | | | | | 27 mg (500 mg) | 1/1/2011 | |
| | Oral | | | | 29.25 mg (500 mg) | 1/1/2011 | |
| | Capsule | | | N/D; N=New, D=Delete | | | |
| | 5 mg | 5/1/2008 | | | | | |
| | 10 mg | 5/1/2008 | | | | | |

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|---------------------------------------|----------------------|----------|-----|------------------------------|--------|-----------|-----|
| MAGNESIUM HYDROXIDE (Age 0-20) | | | | MERCAPTOPURINE | | | |
| | Oral | | | Oral | | | |
| | Suspension | | | Tablet | | | |
| | 2,400 mg/10 mL | 1/1/2011 | | 50 mg | | 2/1/2009 | |
| MAGNESIUM OXIDE (Age 0-20) | | | | METFORMIN | | | |
| | Oral | | | Oral | | | |
| | Tablet | | | Solution | | | |
| | 420 mg | 1/1/2011 | | 500 mg/5 mL | | 2/1/2008 | |
| | 500 mg | 1/1/2011 | | Tablet | | | |
| MECLIZINE | | | | | 500 mg | 2/1/2008 | |
| | Oral | | | 850 mg | | 2/1/2008 | |
| | Tablet | | | 1,000 mg | | 2/1/2008 | |
| | 12.5 mg | 2/1/2008 | | Tablet (24hr) | | | |
| | 25 mg | 2/1/2008 | | 500 mg | | 2/1/2008 | |
| MEDROXYPROGESTERONE | | | | | 750 mg | 2/1/2008 | |
| | Oral | | | METHADONE | | | |
| | Tablet | | | Oral | | | |
| | 2.5 mg | 2/1/2008 | | Concentrate | | | |
| | 5 mg | 2/1/2008 | | 10 mg/mL | | 5/1/2008 | |
| | 10 mg | 2/1/2008 | | Tablet | | | |
| MEGESTROL ACETATE | | | | | 10 mg | 11/1/2009 | |
| | Oral | | | 5 mg | | 11/1/2009 | |
| | Tablet | | | Tablet Sol | | | |
| | 20 mg | 7/1/2009 | | 40 mg | | 8/2/2010 | |
| | 40 mg | 7/1/2009 | | METHENAMINE HIPPURATE | | | |
| MELATONIN | | | | | Oral | | |
| | Oral | | | Tablet | | | |
| | Tablet | | | 1 gram | | 7/1/2009 | |
| | 3 mg | 8/1/2011 | | METHIMAZOLE | | | |
| MELOXICAM | | | | | Oral | | |
| | Oral | | | Tablet | | | |
| | Suspension | | | 10 mg | | 2/1/2008 | |
| | 7.5 mg/5 mL | 2/1/2008 | | 5 mg | | 2/1/2008 | |
| | Tablet | | | METHOCARBAMOL | | | |
| | 7.5 mg | 2/1/2008 | | Oral | | | |
| | 15 mg | 2/1/2008 | | Tablet | | | |
| MERCAPTOPURINE | | | | | 500 mg | 2/1/2008 | |
| | N/D; N=New, D=Delete | | | 750 mg | | 2/1/2008 | |
| | | | | METHOTREXATE | | | |
| | | | | Injection | | | |

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------------------------|-------------------|----------|-----|-------------------------------------------------|----------------------|----------|-----|
| METHOTREXATE | | | | METOCLOPRAMIDE | | | |
| | Injection | | | | Oral | | |
| | Vial | | | | Solution | | |
| | 1 gram | 5/1/2009 | | | 5 mg/5 mL | 2/1/2008 | |
| | 25 mg/mL | 5/1/2009 | | | Tablet | | |
| | 25 mg/mL | 5/1/2009 | | | 10 mg | 2/1/2008 | |
| | Oral | | | | 5 mg | 2/1/2008 | |
| | Tablet | | | METOLAZONE | | | |
| | 2.5 mg | 2/1/2008 | | | Oral | | |
| METHYLDOPA | | | | | Tablet | | |
| | Oral | | | | 10 mg | 2/1/2008 | |
| | Tablet | | | | 2.5 mg | 2/1/2008 | |
| | 250 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | METOPROLOL SUCCINATE/METOPROLOL TARTRATE | | | |
| METHYLDOPA and HYDROCHLOROTHIAZIDE | | | | | Oral | | |
| | Oral | | | | Tablet | | |
| | Tablet | | | | 100 mg | 2/1/2008 | |
| | 250 mg-15 mg | 2/1/2008 | | | 25 mg | 2/1/2008 | |
| | 250 mg-25 mg | 2/1/2008 | | | 50 mg | 2/1/2008 | |
| METHYLPHENIDATE | | | | | Tablet (24hr) | | |
| | Oral | | | | 25 mg | 5/1/2008 | |
| | Tablet | | | | 50 mg | 5/1/2008 | |
| | 5 mg | 2/1/2008 | | | 100 mg | 5/1/2008 | |
| | 10 mg | 2/1/2008 | | | 200 mg | 5/1/2008 | |
| | 20 mg | 2/1/2008 | | METRONIDAZOLE | | | |
| | Tablet ER | | | | Oral | | |
| | 10 mg | 6/1/2010 | | | Tablet | | |
| | 20 mg | 2/1/2008 | | | 250 mg | 2/1/2008 | |
| METHYLPREDNISOLONE | | | | | 500 mg | 2/1/2008 | |
| | Oral | | | MICONAZOLE | | | |
| | Tab Ds Pk | | | | Topical | | |
| | 4 mg | 2/1/2008 | | | Cream | | |
| | Tablet | | | | 2 % | 2/1/2008 | |
| | 4 mg | 2/1/2008 | | | Oinment | | |
| | 8 mg | 2/1/2008 | | | 2 % | 1/1/2011 | |
| METIPRANOLOL | | | | | Vaginal | | |
| | Ophthalmic | | | | Cream | | |
| | Drops | | | | 2 % | 2/1/2008 | |
| | 0.3 % | 2/1/2008 | | | Suppository | | |
| | | | | | 100 mg | 2/1/2008 | |
| METOCLOPRAMIDE | | | | | | | |
| N/D; N=New, D=Delete | | | | | | | |

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------------------------------------|------------------------|----------|-----|----------------------------------------------|--------|----------|-----|
| MICONAZOLE | | | | MORPHINE | | | |
| | Vaginal | | | Oral | | | |
| | Suppository | | | Tablet | | | |
| | 200 mg | 2/1/2008 | | 30 mg | | 1/1/2009 | |
| MIDODRINE | | | | Tablet ER | | | |
| | Oral | | | 15 mg | | 5/1/2008 | |
| | Tablet | | | 30 mg | | 5/1/2008 | |
| | 2.5 mg | 7/1/2009 | | 60 mg | | 5/1/2008 | |
| | 5 mg | 7/1/2009 | | 100 mg | | 5/1/2008 | |
| | 10 mg | 7/1/2009 | | 200 mg | | 5/1/2008 | |
| MINERAL OIL and PETROLATUM | | | | MULTIVITAMINS and IRON (Age 0-20) | | | |
| | Ophthalmic | | | Oral | | | |
| | Ointment | | | Liquid | | | |
| | 15 %-85 % | 1/1/2011 | | | | 1/1/2011 | |
| | 42.5 %-56.8 % | 1/1/2011 | | Tablet | | | |
| MINERAL OIL and PETROLATUM and SODIUM CHLORIDE | | | | | | 1/1/2011 | |
| | Ophthalmic | | | MULTIVITAMINS and MINERALS (Age 0-20) | | | |
| | Ointment | | | Oral | | | |
| | | 1/1/2011 | | Syrup | | | |
| MINOXIDIL | | | | | | 1/1/2011 | |
| | Oral | | | MUPIROCIN | | | |
| | Tablet | | | Topical | | | |
| | 2.5 mg | 2/1/2009 | | Ointment | | | |
| | 10 mg | 2/1/2009 | | 2 % | | 2/1/2008 | |
| MIRTAZAPINE | | | | MYCOPHENOLATE | | | |
| | Oral | | | Oral | | | |
| | Tablet | | | Capsule | | | |
| | 7.5 mg | 2/1/2008 | | 250 mg | | 6/1/2009 | |
| | 15 mg | 2/1/2008 | | Tablet | | | |
| | 30 mg | 2/1/2008 | | 500 mg | | 6/1/2009 | |
| | 45 mg | 2/1/2008 | | NABUMETONE | | | |
| MORPHINE | | | | Oral | | | |
| | Oral | | | Tablet | | | |
| | Solution | | | 500 mg | | 1/1/2010 | |
| | 10 mg/5 mL | 1/1/2009 | | 750 mg | | 1/1/2010 | |
| | 100 mg/5 mL (20 mg/mL) | 1/1/2009 | | NADOLOL | | | |
| | 20 mg/5 mL | 1/1/2009 | | Oral | | | |
| | Tablet | | | Tablet | | | |
| | 15 mg | 1/1/2009 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----|-----------------------------------------------------------------|-----------|-----|-----|-----------------------------------------------------------------|----------|-----|
| | NADOLOL | | | | NEOMYCIN and BACITRACIN and POLYMYXIN and HYDROCORTISONE | | |
| | Oral | | | | Ophthalmic | | |
| | Tablet | | | | Ointment | | |
| | 20 mg | 2/1/2008 | | | 3.5 mg-400 unit-10,000 unit/gram-1 % | 5/1/2008 | |
| | 40 mg | 2/1/2008 | | | | | |
| | 80 mg | 2/1/2008 | | | | | |
| | NALTREXONE | | | | NEOMYCIN and POLYMYXIN and DEXAMETHASONE | | |
| | Oral | | | | Ophthalmic | | |
| | Tablet | | | | Drops Suspension | | |
| | 50 mg | 1/1/2009 | | | 3.5 mg/mL-10,000 unit/mL-0.1 % | 2/1/2008 | |
| | NAPROXEN | | | | Ointment | | |
| | Oral | | | | 3.5 mg/gram-10,000 unit/gram-0.1 % | 2/1/2008 | |
| | Suspension | | | | | | |
| | 125 mg/5 mL | 2/1/2008 | | | NEOMYCIN and POLYMYXIN and HYDROCORTISONE | | |
| | Tablet | | | | Otic | | |
| | 220 mg | 10/1/2009 | | | Drops Suspension | | |
| | 250 mg | 2/1/2008 | | | 3.5 mg/mL-10,000 unit/mL-1 % | 2/1/2008 | |
| | 275 mg | 2/1/2008 | | | Solution | | |
| | 375 mg | 2/1/2008 | | | 3.5 mg/mL-10,000 unit/mL-1 % | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | | | |
| | 550 mg | 2/1/2008 | | | NICOTINE | | |
| | Tablet DR | | | | Buccal | | |
| | 375 mg | 2/1/2008 | | | Gum | | |
| | 500 mg | 2/1/2008 | | | 2 mg | 2/1/2008 | |
| | Tablet ER | | | | 4 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | Transderm | | |
| | NARATRIPTAN | | | | Patch (24hr) | | |
| | Oral | | | | 7 mg/24 hour | 2/1/2008 | |
| | Tablet | | | | 14 mg/24 hour | 2/1/2008 | |
| | 1 mg | 7/1/2011 | | | 15 mg/16 hour | 2/1/2008 | |
| | 2.5 mg | 7/1/2011 | | | 21 mg/24 hour | 2/1/2008 | |
| | NEOMYCIN and BACITRACIN and POLYMYXIN | | | | Patch Step Therapy | | |
| | Topical | | | | 21 mg/24 hour (28)-14 mg/24 hour (14)-7 mg/24 hour (14) | 2/1/2008 | |
| | Ointment | | | | | | |
| | 3.5 mg-400 unit-5,000 unit/gram | 2/1/2008 | | | NIFEDIPINE | | |
| | NEOMYCIN and BACITRACIN and POLYMYXIN and HYDROCORTISONE | | | | Oral | | |
| | Ophthalmic | | | | Capsule | | |
| | Ointment | | | | 10 mg | 2/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----------------------|---------------------|----------|-----|----------------------------------|-----------------------------------------------------------|----------|-----|
| NIFEDIPINE | | | | NITROGLYCERIN | | | |
| Oral | | | | TRANSLING | | | |
| | Capsule | | | | Spray | | |
| | 20 mg | 2/1/2008 | | | 0.4 mg/dose | 2/1/2008 | |
| | TAB ER 24 | | | NORETHINDRONE | | | |
| | 30 mg | 2/1/2008 | | Oral | | | |
| | 60 mg | 2/1/2008 | | Tablet | | | |
| | 90 mg | 2/1/2008 | | | 0.35 mg | 2/1/2008 | |
| | Tablet ER | | | | 5 mg | 8/2/2010 | |
| | 30 mg | 2/1/2008 | | NORETHINDRONE and ETHINYL | | | |
| | 60 mg | 2/1/2008 | | ESTRADIOL | | | |
| | 90 mg | 2/1/2008 | | Oral | | | |
| NITROFURANTOIN | | | | Tablet | | | |
| Oral | | | | | 0.4 mg-35 mcg | 2/1/2008 | |
| | Capsule | | | | 0.5 mg-35 mcg | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | | 0.5 mg-35 mcg (10)/1 mg-35 mcg (11) | 2/1/2008 | |
| NITROFURANTOIN | | | | | 0.5 mg-35 mcg | 2/1/2008 | |
| MACROCRYSTAL | | | | | (7)/0.75 mg-35 mcg | | |
| Oral | | | | | (7)/1 mg-35 mcg (7) | | |
| | Capsule | | | | 0.5 mg-35 mcg (7)/1 mg-35 mcg (9)/0.5 mg-35 mcg (5) | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | | 1 mg-20 mcg | 2/1/2008 | |
| | 50 mg | 2/1/2008 | | | 1 mg-35 mcg | 2/1/2008 | |
| NITROGLYCERIN | | | | | 1.5 mg-30 mcg | 2/1/2008 | |
| Oral | | | | NORETHINDRONE and ETHINYL | | | |
| | Capsule ER | | | ESTRADIOL and IRON | | | |
| | 2.5 mg | 2/1/2008 | | Oral | | | |
| | 6.5 mg | 2/1/2008 | | Tablet | | | |
| | 9 mg | 2/1/2008 | | | 1 mg-20 mcg | 2/1/2008 | |
| | Sublingual | | | | 1 mg-20 mcg (5)/1 mg-30 mcg (7)/1 mg-35 mcg (9)/75 mg (7) | 2/1/2008 | |
| | Sublingual | | | | 1.5 mg-30 mcg | 2/1/2008 | |
| | 0.3 mg | 2/1/2008 | | NORGESTIMATE and ETHINYL | | | |
| | 0.4 mg | 2/1/2008 | | ESTRADIOL | | | |
| | 0.6 mg | 2/1/2008 | | Oral | | | |
| | Transderm | | | Tablet | | | |
| | Oinment | | | | 0.18 mg-35 mcg | 2/1/2008 | |
| | 2 % | 2/1/2008 | | | (7)/0.215 mg-35 mcg | | |
| | Patch (24hr) | | | | (7)/0.25 mg-35 mcg(7)(28) | | |
| | 0.1 mg/hour | 2/1/2008 | | | 0.25 mg-35 mcg | 2/1/2008 | |
| | 0.2 mg/hour | 2/1/2008 | | | | | |
| | 0.4 mg/hour | 2/1/2008 | | | | | |
| | 0.6 mg/hour | 2/1/2008 | | | | | |
| TRANSLING | | | | | | | |
| Spray | | | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----|-----------------------------------------|----------|-----|-----|-------------------------|-----------|-----|
| | NORGESTREL and ETHINYL ESTRADIOL | | | | OFLOXACIN | | |
| | Oral | | | | Ophthalmic | | |
| | Tablet | | | | Drops | | |
| | 0.3 mg-30 mcg | 2/1/2008 | | | 0.3 % | 2/1/2008 | |
| | 0.5 mg-50 mcg | 2/1/2008 | | | | | |
| | NORTRIPTYLIN | | | | OMEPRAZOLE | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Capsule DR | | |
| | 10 mg | 2/1/2008 | | | 10 mg | 4/1/2009 | |
| | 25 mg | 2/1/2008 | | | 20 mg | 4/1/2009 | |
| | 50 mg | 2/1/2008 | | | 40 mg | 4/1/2009 | |
| | 75 mg | 2/1/2008 | | | | | |
| | Solution | | | | ONDANSETRON | | |
| | 10 mg/5 mL | 2/1/2008 | | | Oral | | |
| | | | | | Solution | | |
| | | | | | 4 mg/5 mL | 2/1/2008 | |
| | | | | | Tablet | | |
| | | | | | 4 mg | 2/1/2008 | |
| | | | | | 8 mg | 2/1/2008 | |
| | | | | | 24 mg | 2/1/2008 | |
| | | | | | Tablet Rapdis | | |
| | | | | | 4 mg | 2/1/2008 | |
| | | | | | 8 mg | 2/1/2008 | |
| | NYSTATIN | | | | OSELTAMIVIR | | |
| | Oral | | | | Oral | | |
| | Powder | | | | Capsule | | |
| | 50 million unit | 2/1/2008 | | | 30 mg | 4/29/2009 | |
| | 150 million unit | 2/1/2008 | | | 45 mg | 4/29/2009 | |
| | 500 million unit | 2/1/2008 | | | 75 mg | 4/29/2009 | |
| | Suspension | | | | Suspension | | |
| | 100,000 unit/mL | 2/1/2008 | | | 6 mg/ml | 8/1/2011 | |
| | Tablet | | | | Suspension Recon | | |
| | 500,000 unit | 2/1/2008 | | | 12 mg/mL | 4/29/2009 | |
| | Topical | | | | OXCARBAZEPINE | | |
| | Cream | | | | Oral | | |
| | 100,000 unit/gram | 2/1/2008 | | | Suspension | | |
| | Oinment | | | | 300 mg/5 mL | 2/1/2010 | |
| | 100,000 unit/gram | 2/1/2008 | | | Tablet | | |
| | Powder | | | | 150 mg | 5/1/2008 | |
| | 100,000 unit/gram | 2/1/2008 | | | 300 mg | 5/1/2008 | |
| | Vaginal | | | | 600 mg | 5/1/2008 | |
| | Tablet | | | | OXYBUTYNIN | | |
| | 100,000 unit | 5/1/2008 | | | Oral | | |
| | | | | | Syrup | | |
| | NYSTATIN and TRIAMCINOLONE | | | | | | |
| | Topical | | | | | | |
| | Cream | | | | | | |
| | 100,000 unit/gram-0.1 % | 2/1/2008 | | | | | |
| | Oinment | | | | | | |
| | 100,000 unit/gram-0.1 % | 2/1/2008 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------------------|----------------------|----------|-----|-------------------------------|-----------------------------------------------------------|----------|-----|
| OXYBUTYNIN | | | | PAROMOMYCIN SULFATE | | | |
| | Oral | | | | Oral | | |
| | Syrup | | | | Capsule | | |
| | 5 mg/5 mL | 2/1/2008 | | | 250 mg | 7/1/2009 | |
| | Tablet | | | | | | |
| | 5 mg | 2/1/2008 | | | | | |
| OXYCODONE | | | | PAROXETINE | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Tablet | | |
| | 5 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | Concentrate | | | | 20 mg | 2/1/2008 | |
| | 20 mg/mL | 2/1/2008 | | | 30 mg | 2/1/2008 | |
| | 20 mg/mL (1 mL) | 2/1/2008 | | | 40 mg | 2/1/2008 | |
| | Solution | | | | | | |
| | 5 mg/5 mL | 2/1/2008 | | | | | |
| | Tablet | | | | | | |
| | 5 mg | 2/1/2008 | | | | | |
| | 15 mg | 2/1/2008 | | | | | |
| | 20 mg | 7/1/2011 | | | | | |
| | 30 mg | 2/1/2008 | | | | | |
| | Tablet (12hr) | | | | | | |
| | 10 mg | 2/1/2008 | | | | | |
| | 20 mg | 2/1/2008 | | | | | |
| | 40 mg | 2/1/2008 | | | | | |
| | 80 mg | 2/1/2008 | | | | | |
| OXYCODONE and ACETAMINOPHEN | | | | PEG 3350 | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Soln Recon | | |
| | 5 mg-500 mg | 2/1/2008 | | | 236 gram-22.74 gram- 6.74 gram-5.86 gram- 2.97 gram | 8/1/2009 | |
| | Solution | | | | 240 gram-22.72 gram- 6.72 gram-5.84 gram- 2.98 gram | 8/1/2009 | |
| | 5 mg-325 mg/5 mL | 2/1/2008 | | | | | |
| | Tablet | | | | | | |
| | 5 mg-325 mg | 2/1/2008 | | | | | |
| | 7.5 mg-325 mg | 2/1/2008 | | | | | |
| | 7.5 mg-500 mg | 2/1/2008 | | | | | |
| | 10 mg-325 mg | 2/1/2008 | | | | | |
| | 10 mg-650 mg | 2/1/2008 | | | | | |
| PANTOPRAZOLE | | | | PENICILLIN V POTASSIUM | | | |
| | Oral | | | | Oral | | |
| | Tablet DR | | | | Suspension Recon | | |
| | 20 mg | 7/1/2011 | | | 125 mg/5 mL | 2/1/2008 | |
| | 40 mg | 7/1/2011 | | | 250 mg/5 mL | 2/1/2008 | |
| | | | | | Tablet | | |
| | | | | | 250 mg | 2/1/2008 | |
| | | | | | 500 mg | 2/1/2008 | |
| PERMETHRIN | | | | PENTOXIFYLLINE | | | |
| | | | | | Oral | | |
| | | | | | Tablet ER | | |
| | | | | | 400 mg | 2/1/2008 | |
| PERPHENAZINE | | | | PERMETHRIN | | | |
| | | | | | Topical | | |
| | | | | | Cream | | |
| | | | | | 5 % | 2/1/2008 | |
| | | | | | Liquid | | |
| | | | | | 1 % | 2/1/2008 | |
| | | | | | Oral | | |
| | | | | | Tablet | | |
| | | | | | 2 mg | 2/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|---------------------------------------------------|-------------------------|-----------|-----|-----------------------------------------------------|-------------------------|----------|-----|
| PERPHENAZINE | | | | PHOSPHO SODA SOLUTIONS (Age 0-20) | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Liquid | | |
| | 4 mg | 2/1/2008 | | | 7.2 gram-2.7 gram/15 mL | 1/1/2011 | |
| | 8 mg | 2/1/2008 | | | Solution | | |
| | 16 mg | 2/1/2008 | | | | 1/1/2011 | |
| PHENAZOPYRIDINE | | | | PILOCARPINE | | | |
| | Oral | | | | Ophthalmic | | |
| | Tablet | | | | Drops | | |
| | 100 mg | 11/1/2008 | | | 0.5 % | 2/1/2008 | |
| | 200 mg | 11/1/2008 | | | 1 % | 2/1/2008 | |
| | | | | | 2 % | 2/1/2008 | |
| PHENOBARBITAL | | | | | 3 % | 2/1/2008 | |
| | Oral | | | | 4 % | 2/1/2008 | |
| | Elixir | | | | 6 % | 2/1/2008 | |
| | 20 mg/5 mL | 2/1/2008 | | PILOCARPINE ORAL | | | |
| | Tablet | | | | Oral | | |
| | 15 mg | 2/1/2008 | | | Tablet | | |
| | 16.2 mg | 6/1/2008 | | | 5 mg | 7/1/2009 | |
| | 30 mg | 2/1/2008 | | | 7.5 mg | 8/2/2010 | |
| | 32.4 mg | 2/1/2008 | | PINDOLOL | | | |
| | 60 mg | 2/1/2008 | | | Oral | | |
| | 64.8 mg | 2/1/2008 | | | Tablet | | |
| | 97.2 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| PHENYLEPHRINE and CODEINE and PROMETHAZINE | | | | PIPERONYL BUTOXIDE and PYRETHRINS (Age 0-20) | | | |
| | Oral | | | | Topical | | |
| | Syrup | | | | Liquid | | |
| | 5 mg-10 mg-6.25 mg/5 mL | 5/1/2008 | | | 3 %-0.3 % | 2/1/2008 | |
| | | | | | 4 %-0.33 % | 2/1/2008 | |
| PHENYLEPHRINE and PROMETHAZINE | | | | | Shampoo | | |
| | Oral | | | | 4 %-0.33 % | 2/1/2008 | |
| | Syrup | | | PIROXICAM | | | |
| | 5 mg-6.25 mg/5 mL | 5/1/2008 | | | Oral | | |
| PHENYTOIN | | | | | Capsule | | |
| | Oral | | | | 10 mg | 2/1/2008 | |
| | Capsule | | | | 20 mg | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | POLYMYXIN B and TRIMETHOPRIM | | | |
| | Suspension | | | | | | |
| | 100 mg/4 mL | 2/1/2008 | | | | | |
| | 125 mg/5 mL | 2/1/2008 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-----|-------------------------------------------------------------------------------------------------------------------------------|----------|-----|-----|--------------------------------------------|----------|-----|
| | PREDNISOLONE SODIUM PHOSPHATE | | | | PROCHLORPERAZINE | | |
| | Oral | | | | Rectal | | |
| | Solution | | | | Suppository | | |
| | 5 mg base/5 mL (6.7 mg/5 mL) | 2/1/2008 | | | 25 mg | 2/1/2008 | |
| | PREDNISON | | | | PROMETHAZINE | | |
| | Oral | | | | Oral | | |
| | Concentrate | | | | Syrup | | |
| | 5 mg/mL | 2/1/2008 | | | 6.25 mg/5 mL | 2/1/2008 | |
| | Solution | | | | Tablet | | |
| | 5 mg/5 mL | 2/1/2008 | | | 12.5 mg | 2/1/2008 | |
| | Tab Ds Pk | | | | 25 mg | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | | 50 mg | 2/1/2008 | |
| | Tablet | | | | Rectal | | |
| | 1 mg | 2/1/2008 | | | Suppository | | |
| | 2.5 mg | 2/1/2008 | | | 12.5 mg | 8/2/2010 | |
| | 5 mg | 2/1/2008 | | | PROPAFENONE | | |
| | 10 mg | 2/1/2008 | | | Oral | | |
| | 20 mg | 2/1/2008 | | | Capsule (12hr) | | |
| | 50 mg | 2/1/2008 | | | 225 mg | 7/1/2011 | |
| | PRENATAL VITAMIN-Products packaged as two or more dosage units (i.e. combination packaging) are not a covered service. | | | | 325 mg | 7/1/2011 | |
| | Oral | | | | 425 mg | 7/1/2011 | |
| | | 2/1/2008 | | | Tablet | | |
| | | | | | 150 mg | 5/1/2008 | |
| | | | | | 225 mg | 5/1/2008 | |
| | | | | | 300 mg | 5/1/2008 | |
| | | | | | PROPRANOLOL | | |
| | | | | | Oral | | |
| | | | | | Capsule (24hr) | | |
| | | | | | 60 mg | 5/1/2008 | |
| | | | | | 80 mg | 5/1/2008 | |
| | | | | | 120 mg | 5/1/2008 | |
| | | | | | 160 mg | 5/1/2008 | |
| | | | | | Solution | | |
| | | | | | 20 mg/5 mL | 2/1/2008 | |
| | | | | | 40 mg/5 mL | 2/1/2008 | |
| | | | | | Tablet | | |
| | | | | | 10 mg | 2/1/2008 | |
| | | | | | 20 mg | 2/1/2008 | |
| | | | | | 40 mg | 2/1/2008 | |
| | | | | | 60 mg | 2/1/2008 | |
| | | | | | 80 mg | 2/1/2008 | |
| | | | | | PROPRANOLOL and HYDROCHLOROTHIAZIDE | | |
| | | | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|--------------------------------------------|------------------|----------|-----|-----------------------|-----------|-----------|-----|
| PROPRANOLOL and HYDROCHLOROTHIAZIDE | | | | PYRIDOSTIGMINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 40 mg-25 mg | 2/1/2008 | | | 60 mg | 5/1/2008 | |
| | 80 mg-25 mg | 2/1/2008 | | QUINIDINE | | | |
| PROPYLENE GLYCOL and PEG 400 | | | | | Oral | | |
| | Ophthalmic | | | | Tablet | | |
| | Drops | | | | 200 mg | 2/1/2008 | |
| | 0.3 %-0.4 % | 1/1/2011 | | | 300 mg | 2/1/2008 | |
| PROPYLENE GLYCOL/PEG 400 | | | | | Tablet ER | | |
| | Opth | | | | 300 mg | 2/1/2008 | |
| | Drops Gel | | | RANITIDINE | | | |
| | 0.3-0.4% | 7/1/2011 | | | Oral | | |
| | Drops Liquid Gel | | | | Capsule | | |
| | 0.3-0.4% | 7/1/2011 | | | 150 mg | 2/1/2008 | |
| PROPYLTHIOURACIL | | | | | 300 mg | 2/1/2008 | |
| | Oral | | | | Syrup | | |
| | Tablet | | | | 15 mg/mL | 4/2/2008 | |
| | 50 mg | 2/1/2008 | | | Tablet | | |
| PSEUDOEPHEDRINE | | | | | 150 mg | 2/1/2008 | |
| | Oral | | | | 300 mg | 2/1/2008 | |
| | Syrup | | | RIBAVIRIN | | | |
| | 30 mg/5 mL | 2/1/2008 | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 30 mg | 2/1/2008 | | | 200 mg | 5/1/2008 | |
| | 60 mg | 2/1/2008 | | RIFAMPIN | | | |
| PSYLLIUM (Age 0-20) | | | | | Oral | | |
| | Oral | | | | Capsule | | |
| | Powder | | | | 150 mg | 2/1/2008 | |
| | | 1/1/2011 | | | 300 mg | 2/1/2008 | |
| PSYLLIUM and DEXTROSE (Age 0-20) | | | | RIMANTADINE | | | |
| | Oral | | | | Oral | | |
| | Packet | | | | Tablet | | |
| | | 1/1/2011 | | | 100 mg | 4/29/2009 | |
| | Powder | | | RISPERIDONE | | | |
| | 100% | 1/1/2011 | | | Oral | | |
| PYRIDOSTIGMINE | | | | | Solution | | |
| | | | | | 1 mg/mL | 9/1/2008 | |
| | | | | | Tablet | | |
| | | | | | 0.25 mg | 9/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------|------------|-----------|-----|------------------------------------------------------------------------------|---------------------------------|-----------|-----|
| RISPERIDONE | | | | SERTRALINE | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Concentrate | | |
| | 0.5 mg | 9/1/2008 | | | 20 mg/mL | 2/1/2008 | |
| | 1 mg | 9/1/2008 | | | Tablet | | |
| | 2 mg | 9/1/2008 | | | 25 mg | 2/1/2008 | |
| | 3 mg | 9/1/2008 | | | 50 mg | 2/1/2008 | |
| | 4 mg | 9/1/2008 | | | 100 mg | 2/1/2008 | |
| RIVASTIGMINE | | | | SILVER SULFADIAZINE | | | |
| | Oral | | | | Topical | | |
| | Capsule | | | | Cream | | |
| | 1.5 mg | 10/1/2010 | | | 1 % | 2/1/2008 | |
| | 3 mg | 10/1/2010 | | | | | |
| | 4.5 mg | 10/1/2010 | | | | | |
| | 6 mg | 10/1/2010 | | | | | |
| ROPINIROLE | | | | SIMETHICONE (Age 0-20) | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Drops Suspension | | |
| | 0.25 mg | 7/1/2009 | | | 40 mg/0.6 mL | 2/1/2008 | |
| | 0.5 mg | 7/1/2009 | | | Tablet Chewable | | |
| | 1 mg | 7/1/2009 | | | 80 mg | 2/1/2008 | |
| | 2 mg | 7/1/2009 | | | | | |
| | 3 mg | 7/1/2009 | | | | | |
| | 4 mg | 7/1/2009 | | | | | |
| | 5 mg | 7/1/2009 | | | | | |
| SALSALATE | | | | SIMVASTATIN | | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 500 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| | 750 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| | | | | | 20 mg | 2/1/2008 | |
| | | | | | 40 mg | 2/1/2008 | |
| | | | | | 80 mg | 2/1/2008 | |
| SELEGILINE | | | | SODIUM BICARBONATE | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Tablet | | |
| | 5 mg | 2/1/2008 | | | 325 mg | 1/1/2011 | |
| | Tablet | | | | | | |
| | 5 mg | 2/1/2008 | | | | | |
| SELENIUM SULFIDE | | | | SODIUM CHLORIDE and SODIUM BICARBONATE and POTASSIUM CHLORIDE and PEG | | | |
| | Topical | | | | Oral | | |
| | Suspension | | | | Solution | | |
| | 2.5 % | 2/1/2008 | | | N 420 gms | 10/1/2011 | |
| SERTRALINE | | | | SODIUM FLUORIDE | | | |
| | | | | | Oral | | |
| | | | | | Drops | | |
| | | | | | 0.25 mg fluoride (0.55 mg)/drop | 2/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|----------------------------------------------|-----------------------------|----------|-----|------------------------------------------|----------------------|----------|-----|
| SODIUM FLUORIDE | | | | STAVUDINE | | | |
| Oral | | | | Oral | | | |
| Drops | | | | Capsule | | | |
| | 0.5 mg fluoride (1.1 mg)/mL | 2/1/2008 | | | 15 mg | 2/1/2009 | |
| | 2.2 mg/mL | 2/1/2008 | | | 20 mg | 2/1/2009 | |
| Tablet Chewable | | | | | 30 mg | 2/1/2009 | |
| | 0.25 mg fluoride (0.55 mg) | 2/1/2008 | | | 40 mg | 2/1/2009 | |
| | 0.5 mg fluoride (1.1 mg) | 2/1/2008 | | Solution | | | |
| | 1 mg fluoride (2.2 mg) | 2/1/2008 | | | 1 mg/ml | 7/1/2011 | |
| SODIUM POLYSTYRENE SULFONATE | | | | SUCRALFATE | | | |
| Oral | | | | Oral | | | |
| Powder | | | | Tablet | | | |
| | | 5/1/2009 | | | 1 gram | 5/1/2008 | |
| SORBITOL SOLUTION (Age 0-20) | | | | SULFACETAMIDE and PREDNISOLONE | | | |
| Miscell | | | | Ophthalmic | | | |
| Solution | | | | Drops | | | |
| | 70 % | 1/1/2011 | | | 10 %-0.25 % (0.23 %) | 2/1/2008 | |
| SOTALOL | | | | SULFACETAMIDE SODIUM | | | |
| Oral | | | | Ophthalmic | | | |
| Tablet | | | | Drops | | | |
| | 80 mg | 2/1/2008 | | | 10 % | 2/1/2008 | |
| | 120 mg | 2/1/2008 | | Oinment | | | |
| | 160 mg | 2/1/2008 | | | 10 % | 2/1/2008 | |
| | 240 mg | 2/1/2008 | | SULFAMETHOXAZOLE and TRIMETHOPRIM | | | |
| SPIRONOLACTONE | | | | Oral | | | |
| Oral | | | | Suspension | | | |
| Tablet | | | | | 200 mg-40 mg/5 mL | 2/1/2008 | |
| | 25 mg | 2/1/2008 | | Tablet | | | |
| | 50 mg | 2/1/2008 | | | 400 mg-80 mg | 2/1/2008 | |
| | 100 mg | 2/1/2008 | | | 800 mg-160 mg | 2/1/2008 | |
| SPIRONOLACTONE and HYDROCHLORTHIAZIDE | | | | SULFASALAZINE | | | |
| Oral | | | | Oral | | | |
| Tablet | | | | Tablet | | | |
| | 25 mg-25 mg | 2/1/2008 | | | 500 mg | 2/1/2008 | |
| STAVUDINE | | | | Tablet DR | | | |
| Oral | | | | | 500 mg | 2/1/2008 | |
| | | | | SUMATRIPTAN | | | |
| | | | | Nasal | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|--------------------|-----------------------|----------|-----|---------------------|----------------------|----------|-----|
| SUMATRIPTAN | | | | TERAZOSIN | | | |
| | Nasal | | | | Oral | | |
| | Spray | | | | Capsule | | |
| | 5 mg | 5/1/2010 | | | 1 mg | 2/1/2008 | |
| | 20 mg | 4/1/2009 | | | 2 mg | 2/1/2008 | |
| | Oral | | | | 5 mg | 2/1/2008 | |
| | Tablet | | | | 10 mg | 2/1/2008 | |
| | 25 mg | 2/1/2009 | | TERBUTALINE | | | |
| | 50 mg | 2/1/2009 | | | Oral | | |
| | 100 mg | 2/1/2009 | | | Tablet | | |
| | Sub-Q | | | | 2.5 mg | 2/1/2008 | |
| | Kit-refill | | | | 5 mg | 2/1/2008 | |
| | 4 mg/0.5 mL | 2/1/2009 | | TETRACYCLINE | | | |
| | 6 mg/0.5 mL | 1/1/2009 | | | Oral | | |
| | Pen Inject KIT | | | | Capsule | | |
| | 4 mg/0.5 mL | 2/1/2009 | | | 250 mg | 2/1/2008 | |
| | 6 mg/0.5 mL | 1/1/2009 | | | 500 mg | 2/1/2008 | |
| | Syringe | | | THEOPHYLLINE | | | |
| | 4 mg/0.5 mL | 4/1/2010 | | | Oral | | |
| | 6 mg/0.5 mL | 4/1/2010 | | | Tablet (12hr) | | |
| | Vial | | | | 100 mg | 2/1/2008 | |
| | 4 mg/0.5 mL | 4/1/2009 | | | 200 mg | 2/1/2008 | |
| | 6 mg/0.5 mL | 1/1/2009 | | | 300 mg | 2/1/2008 | |
| | | | | | 450 mg | 2/1/2008 | |
| TACROLIMUS | | | | THIOGUANINE | | | |
| | Oral | | | | Oral | | |
| | Capsule | | | | Tablet | | |
| | 0.5 mg | 9/1/2009 | | | 40 mg | 2/1/2009 | |
| | 1 mg | 9/1/2009 | | THIOTHIXENE | | | |
| | 5 mg | 9/1/2009 | | | Oral | | |
| TAMOXIFEN | | | | | Capsule | | |
| | Oral | | | | 1 mg | 2/1/2008 | |
| | Tablet | | | | 2 mg | 2/1/2008 | |
| | 10 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| | 20 mg | 2/1/2008 | | | 10 mg | 2/1/2008 | |
| TAMSULOSIN | | | | THYROID | | | |
| | Oral | | | | Oral | | |
| | Capsule (24hr) | | | | Tablet | | |
| | 0.4 mg | 6/1/2010 | | | 32.5 mg | 2/1/2008 | |
| TEMAZEPAM | | | | TIMOLOL | | | |
| | Oral | | | | | | |
| | Capsule | | | | | | |
| | 15 mg | 2/1/2008 | | | | | |
| | 30 mg | 2/1/2008 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|--------------------|-------------------------|----------|-----|------------------------|----------------|-----------|-----|
| TIMOLOL | | | | TOPIRAMATE | | | |
| | Ophthalmic | | | | Oral | | |
| | Drops | | | | Tablet | | |
| | 0.25 % | 2/1/2008 | | | 25 mg | 5/1/2009 | |
| | 0.5 % | 2/1/2008 | | | 50 mg | 5/1/2009 | |
| | Oral | | | | 100 mg | 5/1/2009 | |
| | Tablet | | | | 200 mg | 5/1/2009 | |
| | 5 mg | 2/1/2008 | | TORSEMIDE | | | |
| | 10 mg | 2/1/2008 | | | Oral | | |
| | 20 mg | 2/1/2008 | | | Tablet | | |
| TIZANIDINE | | | | | 5 mg | 5/1/2008 | |
| | Oral | | | | 10 mg | 5/1/2008 | |
| | Tablet | | | | 20 mg | 5/1/2008 | |
| | 2 mg | 2/1/2008 | | | 100 mg | 5/1/2008 | |
| | 4 mg | 2/1/2008 | | TRAMADOL | | | |
| TOBRAMYCIN | | | | | Oral | | |
| | Ophthalmic | | | | Tablet | | |
| | Drops | | | | 50 mg | 2/1/2008 | |
| | 0.3 % | 2/1/2008 | | TRANLYCYPROMINE | | | |
| TOLAZAMIDE | | | | | Oral | | |
| | Oral | | | | Tablet | | |
| | Tablet | | | | 10 mg | 10/1/2010 | |
| | 250 mg | 2/1/2008 | | TRAZODONE | | | |
| | 500 mg | 2/1/2008 | | | Oral | | |
| TOLBUTAMIDE | | | | | Tablet | | |
| | Oral | | | | 50 mg | 2/1/2008 | |
| | Tablet | | | | 100 mg | 2/1/2008 | |
| | 500 mg | 2/1/2008 | | | 150 mg | 2/1/2008 | |
| TOLNAFTATE | | | | TRETINOIN | | | |
| | Topical | | | | Oral | | |
| | Cream | | | | Capsule | | |
| | 1 % | 2/1/2008 | | | 10 mg | 7/1/2009 | |
| | Solution | | | | Topical | | |
| | 1 % | 2/1/2008 | | | Cream | | |
| TOPIRAMATE | | | | | 0.025 % | 2/1/2008 | |
| | Oral | | | | 0.05 % | 2/1/2008 | |
| | Capsule Sprinkle | | | | 0.05 % | 2/1/2008 | |
| | 15 mg | 9/1/2009 | | | 0.1 % | 2/1/2008 | |
| | 25 mg | 9/1/2009 | | | Gel | | |
| | Tablet | | | | 0.01 % | 2/1/2008 | |
| | | | | | 0.025 % | 2/1/2008 | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|-------------------------------------------|-------------------|----------|-----|--------------------------|----------------------|-----------|-----|
| TRIAMCINOLONE | | | | TRIMETHOBENZAMIDE | | | |
| | Topical | | | | Oral | | |
| | Cream | | | | Capsule | | |
| | 0.025 % | 2/1/2008 | | | 100 mg | 7/1/2011 | |
| | 0.1 % | 2/1/2008 | | | 250 mg | 7/1/2011 | |
| | 0.5 % | 2/1/2008 | | | 300 mg | 7/1/2011 | |
| | Lotion | | | | TRIMETHOPRIM | | |
| | 0.025 % | 2/1/2008 | | | Oral | | |
| | 0.1 % | 2/1/2008 | | | Tablet | | |
| | Oinment | | | | 100 mg | 2/1/2008 | |
| | 0.025 % | 2/1/2008 | | | TROPICAMIDE | | |
| | 0.05 % | 2/1/2008 | | | Ophthalmic | | |
| | 0.1 % | 2/1/2008 | | | Drops | | |
| | 0.5 % | 2/1/2008 | | | 0.5 % | 2/1/2008 | |
| | | | | | 1 % | 2/1/2008 | |
| TRIAMTERENE and HYDROCHLORTHIAZIDE | | | | | UREA | | |
| | Oral | | | | Topical | | |
| | Capsule | | | | Cream | | |
| | 37.5 mg-25 mg | 2/1/2008 | | | 40 % | 7/1/2009 | |
| | 50 mg-25 mg | 2/1/2008 | | | URSODIOL | | |
| | Tablet | | | | Oral | | |
| | 37.5 mg-25 mg | 2/1/2008 | | | Capsule | | |
| | 75 mg-50 mg | 2/1/2008 | | | 300 mg | 2/1/2008 | |
| | | | | | Tablet | | |
| | | | | | 250 mg | 11/1/2009 | |
| | | | | | 500 mg | 11/1/2009 | |
| TRIFLUOPERAZINE | | | | | VALACYCLOVIR | | |
| | Oral | | | | Oral | | |
| | Tablet | | | | Tablet | | |
| | 1 mg | 2/1/2008 | | | 1,000 mg | 5/1/2011 | |
| | 2 mg | 2/1/2008 | | | 500 mg | 5/1/2011 | |
| | 5 mg | 2/1/2008 | | | VALPROIC ACID | | |
| | 10 mg | 2/1/2008 | | | Oral | | |
| | | | | | Capsule | | |
| | | | | | 250 mg | 2/1/2008 | |
| | | | | | Syrup | | |
| | | | | | 250 mg/5 mL | 2/1/2008 | |
| TRIFLURIDINE | | | | | VENLAFAXINE | | |
| | Ophthalmic | | | | Oral | | |
| | Drops | | | | | | |
| | 1 % | 5/1/2009 | | | | | |
| TRIHEXYPHENIDYL | | | | | | | |
| | Oral | | | | | | |
| | Elixir | | | | | | |
| | 2 mg/5 mL | 2/1/2008 | | | | | |
| | Tablet | | | | | | |
| | 2 mg | 2/1/2008 | | | | | |
| | 5 mg | 2/1/2008 | | | | | |
| TRIMETHOBENZAMIDE | | | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------------------------------|---------|-----------|-----|--------------------|----------|-----------|-----|
| VENLAFAXINE | | | | ZAFIRLUKAST | | | |
| Oral | | | | Oral | | | |
| Capsule (24hr) | | | | Tablet | | | |
| | 37.5 mg | 10/1/2010 | | | 10 mg | 4/1/2011 | |
| | 75 mg | 10/1/2010 | | | 20 mg | 4/1/2011 | |
| | 150 mg | 10/1/2010 | | ZALEPLON | | | |
| Tablet | | | | Oral | | | |
| | 25 mg | 5/1/2008 | | Capsule | | | |
| | 37.5 mg | 5/1/2008 | | | 5 mg | 7/1/2009 | |
| | 50 mg | 5/1/2008 | | | 10 mg | 7/1/2009 | |
| | 75 mg | 5/1/2008 | | ZANAMIVIR | | | |
| | 100 mg | 5/1/2008 | | Inhalation | | | |
| VERAPAMIL | | | | Disk | | | |
| Oral | | | | | 5 mg | 4/29/2009 | |
| Capsule (24hr) Pel | | | | ZIDOVDINE | | | |
| | 120 mg | 2/1/2008 | | Oral | | | |
| | 180 mg | 2/1/2008 | | Capsule | | | |
| | 240 mg | 2/1/2008 | | | 100 mg | 7/1/2011 | |
| Tablet | | | | Syrup | | | |
| | 40 mg | 2/1/2008 | | | 10 mg/mL | 11/1/2009 | |
| | 80 mg | 2/1/2008 | | Tablet | | | |
| | 120 mg | 2/1/2008 | | | 300 mg | 7/1/2011 | |
| Tablet ER | | | | ZOLPIDEM | | | |
| | 120 mg | 2/1/2008 | | Oral | | | |
| | 180 mg | 2/1/2008 | | Tablet | | | |
| | 240 mg | 2/1/2008 | | | 5 mg | 2/1/2008 | |
| VITAMINS A and D (Age 0-20) | | | | | 10 mg | 2/1/2008 | |
| Topical | | | | ZONISAMIDE | | | |
| Cream | | | | Oral | | | |
| | | 1/1/2011 | | Capsule | | | |
| Oinment | | | | | 25 mg | 2/1/2008 | |
| | | 1/1/2011 | | | 50 mg | 2/1/2008 | |
| | | | | | 100 mg | 2/1/2008 | |
| WARFARIN | | | | | | | |
| Oral | | | | | | | |
| Tablet | | | | | | | |
| | 1 mg | 2/1/2008 | | | | | |
| | 2 mg | 2/1/2008 | | | | | |
| | 2.5 mg | 2/1/2008 | | | | | |
| | 3 mg | 2/1/2008 | | | | | |
| | 4 mg | 2/1/2008 | | | | | |
| | 5 mg | 2/1/2008 | | | | | |
| | 6 mg | 2/1/2008 | | | | | |
| | 7.5 mg | 2/1/2008 | | | | | |
| | 10 mg | 2/1/2008 | | | | | |

N/D; N=New, D=Delete

| N/D | STRGTH | Start | End | N/D | STRGTH | Start | End |
|------------|---------------|--------------|------------|------------|---------------|--------------|------------|
|------------|---------------|--------------|------------|------------|---------------|--------------|------------|

1547